R
11
N57
V.184
no.14
cop.2

LIBR
SEP 2
UNIVERSITY

THE BOSTON
Medical and Surgical
JOURNAL

OFFICIAL ORGAN of THE MASSACHUSETTS MEDICAL SOCIETY and of THE NEW ENGLAND SURGICAL SOCIETY

$6 per Annum, 25c per copy
Vol. CLXXXIV, No. 14

THURSDAY, APRIL 7, 1921

Published weekly in Boston, at 126 Massachusetts Avenue

CONTENTS

ADDRESS

ANCIENT EMPIRICISM VS. MODERN EVIDENCE. *By John B. Deaver, M.D.*, Philadelphia.

ORIGINAL ARTICLES

BASAL METABOLISM AS AN INDEX OF TREATMENT IN DISEASES OF THE THYROID. *By Frank H. Lahey, M.D.*, Boston, *and Sara Murray Jordan, Ph.D.*, Boston.

MEDICAL PROGRESS

REPORT ON RECENT PROGRESS IN DERMATOLOGY. *By John T. Bowen, M.D.*, Boston.

EDITORIALS

ENCEPHALITIS WITH EXCITATION.

LICENSING OF PRIVATE HOSPITALS.

For complete table of contents, see first text page.

THE NEXT ANNUAL MEETING OF THE MASSACHUSETTS MEDICAL SOCIETY WILL BE HELD IN BOSTON, TUESDAY, MAY 31, AND WEDNESDAY, JUNE 1, 1921.

The Newest Mayo Clinic Volume

This is by far the largest Mayo Clinic volume ever published. It contains 112 separate articles, totalling 1331 pages, with 496 illustrations, 6 of them in colors. A wide range of subjects is covered. There are 15 articles on the alimentary canal; 29 on the urogenital organs, including 8 on the ductless glands; 2 on the heart; 8 on the blood; 4 on the skin and syphilis; 27 on the head, trunk and extremities; 3 on the nerves; 9 on technic, and 14 general articles.

PARTIAL CONTENTS

Angioma of the stomach
Operability of gastric cancer as determined by x-rays
Cancer of stomach: surgical treatment
Effects of duodenectomy
Routine management of duodenal ulcer
Surgical treatment in bleeding ulcer
Early lesions in the gall-bladder
Jaundice and its significance
Hematogenous infections of the kidney
Recent advance in renal diagnosis
Surgery of the kidney
Stone of the kidney
Surgery of urinary bladder
Treatment of urethral carbuncle
Prostatic diseases
Basal metabolic rates
Thyroid diseases
Organ transplantation
Dangers in blood transfusion
Diagnostic methods in the anemias
Splenectomy in anemia
Syphilis in railroad employees
Tuberculosis of spine
Radium in eye diseases
Mastoiditis
Ankylosis of jaw
Tonsillectomy in myositis and arthritis
Surgery in cysts of thyroglossal tract
Influenza and pneumonia
Antistreptococcus serum in influenza and influenzal pneumonia
Pulmonary suppuration
Fractures as potential deformities
Etiology treatment of acute poliomyelitis
Preparation of prophylactic and autogenous lipovaccines.
Protein sensitization in asthma and hay fever
Experimental surgical shock

Octavo of 1330 pages, illustrated. By WM. J. MAYO, M.D., CHARLES H. MAYO, M.D., and their ASSOCIATES at The Mayo Clinic, Rochester, Minn.
Cloth, $12.00 net.

W. B. SAUNDERS COMPANY — Philadelphia and London

Entered at the Post-office at Boston as second-class matter.



Your Patients Will "Get Well" More Quickly If You Clear Out Their Scybalae

WHEN the intestines are thickly encrusted with dried, hardened feces there isn't much chance for the absorption of nutrient material.

But there *is* ample opportunity for the absorption of products of putrefaction into the circulation.

A large percentage of patients suffering from disorders of nutrition and metabolism are troubled with this condition.

Ordinary cathartics fail completely to loosen and remove these inspissated masses.

Liquid Albolene, by its superior qualities for "mix, penetration and lubrication," facilitates the expulsion of intestinal débris; and thereby promotes absorption of nutritive pabulum.

Try Liquid Albolene—the pure Russian Oil, absolutely bland and non-irritating—on a few of your most obstinate cases of malassimilation, Doctor, and see the improvement manifested in a week or two.

Write for our booklet, "Below The Equator," describing the mode of action of Liquid Albolene.

McKESSON & ROBBINS, INC.

MANUFACTURING CHEMISTS

Established 1833 **NEW YORK CITY**

HAY FEVER MEMORANDA

EARLY SPRING TYPE. Patients whose hay fever develops in late March, April or beginning of May should be tested with pollens of early flowering trees.

LATE SPRING TYPE. Patients whose hay fever develops in the latter part of May, or during June, should be tested with the pollens of the chief grasses, such as orchard grass, timothy, red top—and certain early flowering weeds.

ARLCO-POLLEN EXTRACTS for Cutaneous Tests and Treatment cover early and late spring, also summer and autumn.

Literature and list of Pollens on request.

THE ARLINGTON CHEMICAL CO.

Yonkers, New York

ACRIFLAVINE - HEYL
PROFLAVINE - HEYL

Have been accepted by the Council on Pharmacy and Chemistry of the A.M.A. for inclusion in New and Non-Official Remedies.

Used in the treatment of Gonorrhoea for both acute and chronic conditions.

Clinical reports show that both Acriflavine-Heyl and Proflavine-Heyl have proven valuable in the treatment of wound infections with a slight advantage in favor of Acriflavine-Heyl.

Recent work proves their value in Dermatology.

VEN-ACRIFLAVINE for intravenous injection is a 1-3000 solution of Acriflavine-Heyl specially prepared in normal saline solution. Sold in 5 cc. ampoules, six in a carton.

Highly diffusible and less irritating than the organic silver salts, their potency is increased in the presence of the serum.

Write for literature and price lists to

HEYL LABORATORIES Inc., 146th St. and Park Ave., New York City

TESTOGAN
For Men

THELYGAN
For Women

Formula of Dr. Iwan Bloch

After seven years' clinical experience these products stand as proven specifics.

INDICATED IN SEXUAL IMPOTENCE AND INSUFFICIENCY OF THE SEXUAL HORMONES

They contain SEXUAL HORMONES, i. e., the hormones of the reproductive glands and of the glands of internal secretion.

Special Indications for Testogan:

Sexual infantilism and eunuchoidism in the male. Impotence and sexual weakness. Climacterium virile. Neurasthenia, hypochondria.

Special Indications for Thelygan:

Infantile sterility. Underdeveloped mammae, etc. Frigidity. Sexual disturbances in obesity and other metabolic disorders. Climacteric symptoms, amenorrhea, neurasthenia, hypochondria, dysmenorrhea.

Furnished in TABLETS, for internal use, and in AMPOULES, for intragluteal injection.
Prices: $2.00 per box of 40 tablets; $3.00 per box of 20 ampoules

EXTENSIVE LITERATURE ON REQUEST

CAVENDISH CHEMICAL CORPORATION

220 West 43rd Street — *Sole Agents* *Established 1905* — New York

NOTICE

SHERMAN'S VACCINES

ARE NOW SUPPLIED IN A NEW 10 MIL. (C. C.) CONTAINER

10 Mil. (c. c.)

Twenty Preparations Beyond the experimental stage.

Millions of doses have been administered.

This package has many superior features which assure asepsis, prevent leakage and facilitate the removal of contents. It is constructed on the well known Sherman principle.

The vial is amply strong which prevents breakage so frequent with shell vials.

We are exclusive and pioneer producers of Bacterial Vaccines. Originators of the aseptic bulk package. Pioneer in elucidation, experimentation and clinical demonstration.

The largest producers of Stock and Autogenous Bacterial Vaccines.

MANUFACTURER OF BACTERIAL VACCINES

G. H. SHERMAN M.D.
Detroit, Mich. U.S.A.

"Sherman's Vaccines are dependable Antigens"

"E.F.M." Perfect Catgut

The Catgut with a reputation for

Sterility—Suppleness Strength

The "E. F. M." Catgut is found in the operating rooms of most of the hospitals in New England and is used by many of the most prominent surgeons. They realize that the "E. F. M." is the last word in catgut production.

E. F. Mahady Company

671 Boylstont Sreet BOSTON, MASS.

Miscellany.

VENEREAL DISEASE CONFERENCE.

A conference attended by representatives of the United States Public Health Service, State Health Departments, and of a clinic of national repute was held in Washington, D. C., on January 12, 1921, to discuss the work of venereal disease control and to propose plans for further development. The following resolutions were adopted:—

Resolved: 1. That the educational work of the venereal disease control and social hygiene movements has been sufficiently effective in developing a sympathetic and supportive public sentiment to warrant its continuation.

2. That an effort should now be made to evaluate the educational work in as exact social, psychologic, and medical terms as possible.

Resolved: That the administration of public health measures relating to venereal disease control should apply equally to both sexes.

Resolved: That health officers should give preference to education and persuasion before having recourse to legal process in connection with enforcing laws and regulations relating to venereal disease control.

Resolved: That while the United States Public Health Service and the State Boards of Health recognize their public duty to see that every victim of a contagious disease received adequate treatment for his own and the public's safety, they have no intention of supplanting effective private effort in this field.

Resolved: That there is urgent need for all physicians to recognize their responsibility to the community and the patient, in the control and the treatment of venereal disease, by themselves raising the standard of treatment. This implies that a physician who is unfamiliar with, or unprepared to employ modern methods in the management of these diseases, should not accept such cases for treatment but should refer them to some private or public physician who is properly equipped.

Resolved: That the venereal disease control movement cannot reach full effectiveness without the intelligent and sympathetic coöperation of the medical profession. This coöperation can best be obtained by:

1. The rapid extension of teaching facilities for medical students so that knowledge of the

(Continued on page vi.)

SANBORN

Blood Pressure Outfit

Thorough accuracy, extreme durability, quick and dependable determination of systolic, diastolic and pulse pressures—these are guaranteed the user of the Sanborn Outfit.

Approved by the U. S. Army and Navy, insurance companies and thousands of physicians throughout the country.

Aneroid gage; washable armband, 42 inches long; Ceylon rubber bag, 5 × 9½ inches, reinforced by durable fabric; specially made release valve; essential parts double strength; packed in case of selected leather.

PRICE — $17.50

May we not send you an outfit for trial in your practice? No advance payment required. Just send your request on your business stationery.

Made, guaranteed and sold direct to physicians by

Sanborn Company Makers of Scientific Instruments

1048 Commonwealth Ave., Boston, 47, Mass.

SURGEONS and PHYSICIANS SUPPLY CO.

Silver Salvarsan

Manufactured by

H. A. Metz Laboratories

IMMEDIATE DELIVERY

0.05 gram	.....	$ .75	per ampule
0.1 "	.....	1.00	" "
0.15 "	.....	1.25	" "
0.2 "	.....	1.50	" "
0.25 "	.....	1.75	" "
0.3 "	.....	2.00	" "

10% discount in 10 ampule lots

Sudbury Building, Boston, Sudbury St. Mass.

Tel. Hay. 1756

MANY ADVANTAGES AS A DIET FOR SURGICAL CASES

THE ORIGINAL

Promotes Convalescence

Avoid imitations

Samples prepaid upon request

HORLICK'S MALTED MILK CO.

RACINE, WIS.

Probilin Pills

BILIARY AND HEPATIC

Fifteen years of extensive universal use of Probilin Pills have demonstrated that by early diagnosis and logical medical treatment a great many gall stone cases can be prevented from reaching the stage of indispensable surgical intervention.

Indispensable operation.

That's the point.

So many are performed that are not.

Probilin Pills literature on request.

MADE IN U.S.A.

FURNISHED SIXTY IN AMBER VIAL WITH GREEN LABEL AND RED SEAL AT BEFORE-THE-WAR PRICE.

SCHERING & GLATZ, INC. NEW YORK

(Continued from page iv.)

medical, social, and public health aspects of these diseases may be taught by actual contact with patients in the clinic under the direction of qualified teachers.

2. By making available to all physicians by means of clinics, lectures, demonstrations, and institutes, the most recent developments in medical and social knowledge of the venereal diseases.

3. By the development of state diagnostic facilities for the use of the practising physician.

Plates for the Relief of Flat Foot

$3.00 A PAIR, INCLUDING FITTING

Made to Order from Casts and Directions

From strong, non-corrosive material. Directions for making casts on application

INSTRUMENT REPAIRING A SPECIALTY

CODMAN & SHURTLEFF, Incorporated

Mfrs. of Superior Surgical Insts. Established 1838

120 Boylston St. Boston, Mass.

95 YEARS

experience has proven

WINSOL

A GUARANTEE OF QUALITY

Cerizane Balsam

to be the most resultful vegetable expectorant. Contains no opium, narcotics or chloroform.

Can always be found at any Winsol Store, or any druggist can get it through his jobber.

B. O. & G. C. WILSON, Inc.

(Established 1845)

BOSTON, - MASS.

Resolved: That the development and maintenance of a competent medical personnel under executive leadership in the field of venereal disease control demands training, whose duration and character is varied with the work for which preparation is sought; from six months, for the routine performance of the technical work of treatment, to three or more years for the preparation of men who are to assume the full medical and administrative responsibilities of the expert. For the purpose of such training, special post-graduate courses in association with medical schools and teaching centers providing liberal and accessible clinical material are essential. The development of such schools should be encouraged and active participation of their graduates in this work should be sought. The maintenance of an efficient personnel further requires the provision of inspiration and incentive to individual development and reward for initiative comprised in (a) adequate material equipment for laboratories and clinics; (b) adequate technical and medical assistance for the handling of routine work; (c) sufficient freedom from routine and provision of funds to make possible the conduct of research; (d) much more generous provision for the salary of personnel than is now the rule.

The continuance of public parsimony in this field will ultimately divert from the public service, into more generously rewarded lines of activity, those men whose training and capacity can redeem the work from mediocrity and the movement from futility.

Resolved: That the obligation of a public or private agency for the treatment of venereal disease does not end with the mere overcoming of a group of symptoms or the temporary control of contagiousness, but should extend throughout the course of the disease. The fact that a person has a venereal disease should not, as such, act as a bar to his admission to any hospital or institution receiving public funds. The aim of all agencies for the care of venereal diseases should be to trace out infected individuals; to carry treatment to the point of cure or arrest; to accumulate a body of records for intelligent control of the individual case, and to further scientific research; to maintain a follow-up system and to provide special diagnostic facilities; and to provide careful and repeated observation and expert advice for the individual patient throughout life.

LOESER'S INTRAVENOUS SOLUTIONS

Sodium Iodide

A sterile solution in sealed glass ampoules; 20 c.c. contain 2 gm. (31 grains) Sodium Iodide U. S. P.

The beneficial effects of iodides, especially of sodium iodide, are enhanced and rendered almost certain by introducing the drug directly into the blood stream.

In a broad class of cases, including Asthma, Rheumatism, Orchitis, late Syphilis, Gonorrheal secondaries, arteriosclerosis, etc., sodium iodide intravenously injected secures results.

Loeser's Intravenous Solution of Sodium Iodide is properly made in a scientific laboratory by scrupulously exact methods.

It is stable, uniform, NON-IRRITATING—SAFE.

Intravenous administration of drugs is the coming therapy.

Its results are quicker, more certain.

The method proves popular with the patient. It gains prestige for the physician.

It has passed the introductory stage—is rapidly coming into general use. Loeser's Intravenous Solutions are Standard.

Clinical reports, reprints, list of products, price list, directions for use, etc., sent to any physician on request.

Arsenic and Mercury
Sodium Iodide
Salicylate and Iodide
Sodium Salicylate
Mercury Bichloride
Mercury Oxycyanide
Quinine Dihydrochloride
Hexamethylenamine
Calcium Iodide

New York Intravenous Laboratory
112 West 21st Street, New York

RADIUM

STANDARD CHEMICAL CO.

RADIUM of highest purity in any quantity.

Patented glazed plaques
for superficial conditions.

Tube and Needle Applicators
for deep therapy.

Apparatus for radium emanation
installed by our Department of Physics.

All our applicators and apparatus adopted after having been proven therapeutically practicable.

U. S. Bureau of Standards Certificate.

Our Departments of Physics and Medicine give instruction in the physics and therapeutic application of Radium.

RADIUM CHEMICAL CO.

PITTSBURGH, PA.

BOSTON CHICAGO SAN FRANCISCO NEW YORK

We are producing an

ARSPHENAMINE

AND A

NEOARSPHENAMINE

than which no better products are made anywhere in the world

Recent studies carried out in the U. S. P. H. S. Laboratories at Washington, comparing the toxicity of the American and Foreign arsphenamines and neoarsphenamines, demonstrate that the American products exhibit a lower toxicity.

Analogous studies made in our own laboratories give similar results.

Our neoarsphenamine is rapidly soluble and may be administered in 5 c.c. of water. It is splendidly tolerated and produces admirable clinical and serological results.

DERMATOLOGICAL RESEARCH LABORATORIES

1720-22 Lombard Street, and 1713-15-17-19-21-23 Naudain Street.

Philadelphia

DRL

Massachusetts General Hospital Pay Consultation Clinic

To aid in diagnosis in cases requiring the advice of several specialists, members of the staff of the Massachusetts General Hospital will hold clinics at the hospital Tuesday and Friday afternoons at 1.30 P.M.

Physicians desiring to make use of the clinic should refer their cases by letter addressed to the Consultation Clinic, giving a brief history. The object being to establish a diagnosis, no treatment will be given, nor will patients be admitted to the hospital except to complete an examination which cannot be completed in the limited time of the clinic.

Patients are referred back to the doctor who sends them to us. Several visits may be necessary to establish a diagnosis, and when ordered by the clinic staff, no additional charge will be made for such visits. Upon completion of the examination a report will be sent to the doctor who refers the case.

No appointment is necessary.

No treatment given.

Hours: Tuesdays and Fridays, 1.30 P.M. to 2 P.M.

A letter from a doctor is required in every case.

Consultation	**$10.00**
X-Ray	**3.00 to 10.00**
Laboratory examination	**2.00**
Cystoscopy	**3.00**
Electrocardiogram	**10.00**

Departments represented:

Medicine	**Surgery**
Genito-urinary	**Orthopedics**
Syphilis	**Skin Diseases**
Pediatrics	**Neurology**
Nose and Throat	**Eye and Ear**

F. A. WASHBURN, Sec'y,
General Executive Committee.

EVERY PHYSICIAN

In general practice should have a BURDICK ELECTRIC LIGHT APPLICATOR for local and general applications. It will produce

DEEP
EFFECTIVE
HYPERAEMIA
INCREASED
LEUCOCYTOSIS
INCREASED
VITAL
RESISTANCE

Useful for the relief of inflamation and congestion, to promote quick healing of wounds, relieve pain, etc.

WRITE FOR DISCRIPTIVE LITERATURE

CLAPP ANDERSON CO.

Specialists in high quality X-Ray and Electro-Medical Apparatus

120 BOYLSTON STREET, BOSTON, MASS.

Prescribe *"By Far The Best"*

Burkhardt BREWING COMPANY

Products for Medicinal Use

Burkhardt's "THE GENUINE"

for medicinal purposes affords "By Far The Best" in Malt beverages. A product of nearly 75 years' experience. First brewed in 1847. We know only the choicest ingredients are used. "By Far The Best" the world's markets afford. This long established reputation for QUALITY suggests confidence in this present day product. Prescribe it! You will make no mistake.

AS YOU PRESCRIBE
MALT BEVERAGES
MALT EXTRACT

Prescribe by Name Burkhardt's "THE GENUINE"

Burkhardt Brewing Co.
Boston 20, Mass.
Telephones
Roxbury 4456, 4457, 4458

TEROGON

A New Antigonorrheal for Internal Administration

COMPOSITION:

Extract of Pichi-Pichi	75.
Extract of Kawa-Kawa	14.5
Lecithin	5.5
Ethereal Oil of Chamomile	0.01

Used by Dr. ADOLF SCHUFTAN with good results in over 200 cases

Send for
"9 Reasons"
why Terogon should be used in preference to any other internal treatment for gonorrhea. Mailed free on request.

Special introductory price to physicians:
$1.00 per box of 60 tablets

CAVENDISH CHEMICAL COMPANY

220 WEST 43rd STREET, NEW YORK

We are the sole American Agents for Yohimbin (Henning)
Prices: In tubes of 10, $1.00; in boxes of 40, $3.00

DESIDERATA

A lactic culture, to properly perform its function, must be rapidly viable, easy of administration and harmless in any quantity, even though the patient be a delicate infant. A convenient form and simplicity of product are minor advantages which are appreciated by the physician.

When prescribing a culture of Bacillus Bulgaricus give

B. B. CULTURE

a trial. The results will be most gratifying.

B. B. CULTURE LABORATORY
INCORPORATED
YONKERS, NEW YORK

LIBERAL DOSAGE—MINIMUM EXPENSE

In specializing on Bacillus Bulgaricus Cultures under the scientific guidance of Professor Metchnikoff since 1906 and by constant co-operation and collaboration with eminent medical practitioners, we have arrived at the conviction that **adequate dosage** is one of the most important factors in the effective application of the Bacillus Bulgaricus to therapeutic purposes. Seven c.c. tubes have brought satisfactory results in cases where long-continued use of tablets showed no improvement whatsoever, and the administration of one ounce or two ounce doses, once or twice daily, gave surprisingly rapid results where the 7 c.c. doses had proved of little benefit, even when administered three to six times daily.

Acting on this scientific conclusion, we have worked out the problem of LIBERAL DOSAGE—MINIMUM EXPENSE. We have always maintained, and still steadfastly maintain, that it is unscientific and even dangerous to dispense culture media in bulk. Our reason for this has been clearly set forth for years on page 6 of our Physician's List. We have, therefore, introduced the "small drink" dose in individual, one-dose bottles.

THE BIG THREE

(ACTUAL SIZE)

"PINEAPPLE CULTURE, METCHNIKOFF." A two-ounce dose of Culture A in each bottle. Suitable for all cases of auto-intoxication, except Diabetes, or where fruit juices are counter-indicated. Price per case of one dozen bottles, delivered to patient's address, anywhere east of Mississippi River, $2.00.



"WHEY CULTURE BACILLUS BULGARICUS, METCHNIKOFF." A one and one-half ounce dose in each bottle. Suitable for colon injection and all cases where a fruit juice medium is not desired. Price per case of one dozen bottles, delivered to patient's address, anywhere east of Mississippi River, $2.25.

"BULGARICUS BOUILLON, METCHNIKOFF." For Diabetics. A one and one-half ounce dose of Culture D in each bottle. The most valuable therapeutic agent for the treatment of Diabetes known to medical science. Price per case of one dozen bottles, delivered to patient's address, anywhere east of Mississippi River, $2.25.

All the above products can be obtained at the leading druggists or will be delivered at patients' address without extra charge.

Literature and Samples on Request

FRANCO-AMERICAN FERMENT COMPANY

225-7 SIXTH AVENUE, NEW YORK

Yeast treatment successful in controlling constipation

THE thorough-going trial of yeast as a therapeutic agent carried on at Jefferson Medical College, the Philadelphia General Hospital and the Roosevelt Hospital in New York indicated clearly the great value of yeast in the control and cure of constipation.

Ten cases were treated. Only one refused to yield — a case complicated by subacute appendicitis with high intestinal stasis.

Of the nine cases that yielded readily to yeast treatment the following are typical:

Slight constipation for a few weeks; treatment, 2 cakes daily before meals for a month; results, regular movements from the start of yeast. Chronic constipation for a year or more receiving 3 compound cathartic pills daily; treatment, 3 cakes yeast daily before meals; results, movements became regular without other medication.

Another case involved constipation for several years with periodic headache. The treatment of 3 cakes of yeast daily before meals resulted in regular movements within 4 or 5 days. In addition to the relief from the specific symptoms in question the general physical condition improved.

The final conclusion of those who treated and observed these cases is: "It seems to us that yeast is especially indicated in chronic non-obstructive bowel conditions and contra-indicated in the acute conditions."

In these tests Fleischmann's Yeast was used — the same that is used by countless housewives in making bread. This yeast is of the species Saccharomyces Cerevisiæ, scientifically cultured, and kept free from contamination by wild yeast or other impurity.

Usual dosage one cake yeast t.i.d. before or between meals, administered in a suspension of water, fruit-juices or milk.

Fleischmann's Yeast may be secured fresh, daily, from all grocers. If the physician prefers, he may write to The Fleischmann Company in the nearest large city and a supply will be mailed direct on the days wanted.

THE FLEISCHMANN COMPANY
701 Washington Street, New York, N. Y.

TWENTY years ago Parke, Davis & Co. introduced to the medical profession the active principle of the suprarenal gland—Adrenalin.

Little was known at that time concerning its physiologic action and therapeutic application. Today, after years of laboratory research and clinical experimentation, Adrenalin holds a foremost place among the standbys of the materia medica.

For the relief of the paroxysm of asthma, for the control of hemorrhage, and in the treatment of shock and collapse, Adrenalin is the first thought of the therapeutist. In organotherapy it has certain special indications, and as a synergist to local anesthetics it has done much toward bringing local anesthesia technic to its present high degree of perfection.

PARKE, DAVIS & CO.

CANCER ODORS

—and the malodorous dejecta of typhoid and other intestinal cases come under the same category—

can be made materially less prejudicial to the nurse's health if you permit her to insert a tuft of "Nazeptic Wool" in each nostril before she dresses the cancer or empties the used vessel. "Nazeptic Wool" is absorbent cotton medicated with phenol, eucalyptol, methyl-salicylate and menthol; such odors, when filtered through the "Wool" during inhalation, are materially antisepticized. Samples are available.

SHARP & DOHME of Baltimore

Have You Seen the Latest Material from the FOREIGN Journals?

VALUABLE FOREIGN MATERIAL LOST

A vast amount of invaluable information appearing every month in the foreign medical journals is lost to American physicians because they are unable to secure these journals, or else because of their inability to translate more than one or two languages.

The Library Department of the American Institute of Medicine prepares, each month, a large number of abstracts of articles taken from foreign journals. Translations are made from the following languages:

TRANSLATIONS AND ABSTRACTS MADE FOR YOU

FRENCH	SPANISH
GERMAN	SCANDINAVIAN
ITALIAN	RUSSIAN

By making use of the services of the Library Department any physician can easily keep in touch with the best material appearing in the foreign journals. This Department also furnishes complete index and abstracts of all important articles appearing in the American, Canadian and British journals.

AN EASY METHOD OF KEEPING UP-TO-DATE

The complete "Survey" is carefully indexed and classified according to subject, printed in loose leaf form and furnished in convenient library size ring binder. Abstracts on all medical subjects may be had, or if desired, only those abstracts relating to one particular subject will be sent.

SPECIMEN ABSTRACTS SENT GRATIS

Physicians who would like to know about the work of the Library Department of the American Institute of Medicine are requested to fill out the attached coupon. Explanatory literature and specimen index and abstracts will be sent by return mail.

American Institute of Medicine

13 East 47th Street, NEW YORK CITY

COUPON

AMERICAN INSTITUTE OF MEDICINE
13 East 47th Street
New York City

Date .....

Please send literature containing detailed information about the Library Department —also specimen abstracts and index. I am especially interested in .....
(Name of subject)

Name .....
(Please print)

Street .....

Town .....

State .....

(B.M.S.J. 4-7-21)

BUTTER *vs.* SUBSTITUTES

The comparative price of butter *vs.* oleomargarines mislead the public into believing that substitutes are much more economical. While butterfat is rich in the fat soluble vitamines, plant oils contain practically none.

Those families who use butter substitutes should use sufficient *milk* to supply the necessary growth promoting substances. It is interesting to note the results of a recent experiment conducted by a Jewish Orphan Asylum in Rochester, N. Y.—a small asylum containing ten children. Believing that they could economize by substituting oleomargarine for creamery butter, a six months' test was conducted, with the result that every child lost weight. Butter was again introduced and the children immediately began to grow.

Butter is a concentrated, vital food.

Our butter is fresh-churned from carefully ripened cream.

We make daily home delivery throughout Boston and Suburbs, express and baggage shipments to other points.

D. WHITING & SONS, Boston

The Management of an Infant's Diet

Constipation

Infants that have a fat intolerance as a result of being fed upon mixtures containing a high percentage of cow's milk fat may develop a condition of constipation of a most pronounced type. This appears, at first, to be most difficult to correct; yet a very simple adjustment of the diet will bring prompt relief. The proper procedure is to remove all of the cream from the milk to be used in preparing the diet and add an easily assimilable carbohydrate. This carbohydrate element should be free from starch and one that has a high point of assimilation, for it is important that a relatively high percentage be used in order to compensate for the heat and energy lost by removing the fat.

Mellin's Food

contains the carbohydrate—maltose—which answers the purpose, for maltose is utilized in larger amounts than any other carbohydrate. Successful results may therefore be obtained by preparing the modification with skimmed milk and at least four level tablespoonfuls of Mellin's Food to each pint of the food mixture.

Further details are given in a pamphlet which physicians may obtain upon request.

Mellin's Food Company
Boston, Mass.

The Boston Medical and Surgical Journal

TABLE OF CONTENTS

April 7, 1921

ADDRESS

ANCIENT EMPIRICISM VS. MODERN EVIDENCE. *By John B. Deaver, M.D.*, Philadelphia. 345

ORIGINAL ARTICLE

BASAL METABOLISM AS AN INDEX OF TREATMENT IN DISEASES OF THE THYROID. *By Frank H. Lahey, M.D.*, Boston, *and Sara Murray Jordan, Ph.D.*, Boston. 348

MEDICAL PROGRESS

REPORT ON RECENT PROGRESS IN DERMATOLOGY. *By John T. Bowen, M.D.*, Boston. 358

BOOK REVIEW

Tropical Ophthalmology. By Robert Henry Elliot, M.D., B.S., F.R.C.S. 362

EDITORIALS

ENCEPHALITIS WITH EXCITATION. 363

LICENSING OF PRIVATE HOSPITALS. 363

MEDICAL NOTES. 364

CORRESPONDENCE

ACTION TAKEN BY THE PROFESSIONAL GUILD OF KING'S COUNTY. *John P. Davin*, New York, N. Y. 365

THE DANGER OF TYPHUS. *S. G. Pavlo, M.D.*, Malden, Mass. 366

SHARON SANATORIUM NOT A STATE INSTITUTION. *Vincent Y. Bowditch, M.D.* 366

MISCELLANY

SOCIETY NOTICES. 366

Address

ANCIENT EMPIRICISM VS. MODERN EVIDENCE.*

BY JOHN B. DEAVER, M.D., PHILADELPHIA.

IT is a pleasure to me to be with you on this occasion which to me represents an expression of the spirit of fellowship and coöperation among the members of our professional fellow-citizens of this state. At the same time, I would like to express the hope that in the near future there may be found a way for you to join forces with the state branch of the national organization, of which I have the honor of being President-elect, so that there may be that one hundred per cent. coöperation so essential to improving and maintaining the standards of our calling.

I cannot help yielding to a natural and compelling impulse to preface what I have to say to you this evening by extending to the future vice-president the congratulations of this assembly, both in its professional capacity and its capacity as loyal Americans on the complete and signal victory experienced at the polls. There is every reason, not so much to hope as to believe, that the prospective change in administration must result in the improved conditions so anxiously awaited by the country at large. This is neither the time nor the place to enter into a discussion of the disadvantages under which this country has been traveling for the past few years. But it is both proper and fitting to express gratification at the turn of events and to pledge our enthusiastic support to the new heads of our national government. This pledge is all the more significant because it is given also to a scion of your state and because of the important rôle played by Massachusetts in the history of these United States. It is but maintaining its tradition in giving one of its sons the opportunity of helping to add another illustrious page to that glorious history of Massachusetts. Not only to the Vice-President-elect, but also to the Governor-elect of the state, it gives me pleasure to offer my best wishes for his success in continuing the policy which led to the choice of Governor Coolidge for the high office to which he has just been elected.

But not alone in the political history of our country does Massachusetts stand supreme, in medical annals also she occupies an enviable position. If there were no other achievement to her credit than that within her borders were first published the experiments that resulted in the boon of anesthesia, it were enough

* Read before the meeting of the Clinical and Surgical Association of Massachusetts, Boston, Nov. 9, 1920.

to emblazon the name of Massachusetts in bold letters on the honor roll of our guild. Not content with this signal victory, she has enriched and is still enriching our profession with a galaxy of names which represent and will always stand for the highest ideals and accomplishment. The list of epoch-making contributions furnished to the medical sciences by physicians of Massachusetts, and more especially of Boston, is no mean one. Many of these were pioneers in their special fields. In the early days you gave us the Warrens, Keep, the Bigelows, Hare, Fischer, the inventor of the cystoscope; then the beloved poet-physician, Oliver Wendell Holmes, whose discovery of the cause of puerperal fever paved the way for the protection of the mother at her critical time, and has spared countless lives; and Reginald Heber Fitz, the revealer of the true malefactor in the disease which bears the name he gave it, and who also shed light on the rôle of the pancreas in certain, then obscure, abdominal conditions; Maurice Richardson, whose acumen and forethought have done much to place surgery in the estimable position it now holds. And thus I could go on. The list is a large one and, no doubt, more familiar to you than to me. You and we are fortunate in having still with us, men of Massachusetts who continue to form the Hub from which new ideas and new methods continue to radiate to all parts of the globe, and who are keeping alive the traditions of pioneer New England.

There are certain topics that never lose their interest, and among these I believe the early efforts in surgery retain a remarkable fascination. The art of surgery is as old as history itself, and in the course of time it has come to embrace every part of the human organism. In the chronicles of the ages surgery apparently antedates medicine, the latter being more intimately bound up with religion and with priestly functions, and at all times the former has carried with it a greater responsibility than the practice of the healing art as such. We see this in the Babylonian code of laws in which, we are told, the blame for failure of an operation rested entirely upon the surgeon for his lack of judgment in taking the risk. The punishment for bad surgery was made to fit the crime, and also the patient. If an unsuccessful operation was done on a slave, that is, if the patient—a slave—lost an eye, for example, the bungling surgeon was obliged to pay half the value of the slave; if the patient died, a new slave was required of the operator. If the victom was a free man, the penalty in either contingency—the loss of an eye, or death—consisted in cutting off the hand of the operator. Not an unwise precautionary measure, fortunately (or unfortunately) not applicable today! The early Roman laws were equally severe on the erring surgeon. If, when called upon to bleed a patient, he drew too much blood, he was obliged to pay a liberal indemnity to his patient, and if the latter died, the surgeon's life was at the mercy of the family of the defunct. If the patient happened to be a slave, restitution of a slave of equal value was required. The law, however, dealt more leniently with the physician, although failure to cure was attended with loss of his fee. But, according to the naïve statement of an ancient document the physician is better able to hide his mistakes than is the surgeon, and so we may presume that the doctors of those days collected their fees pretty regularly.

The surgeon of other days likewise labored under the disadvantage of occupying a lower social and professional status than the physician. Manual treatment of any kind being considered beneath the dignity of the medical practitioner, the surgeon was called upon to act as hands to the latter, so that surgery became a sort of trade followed principally by barbers. It is only within comparatively recent times that the two professions have been put on an equal footing. The attainment was not a sudden nor an easy one. It has been a slow, steady evolution not unattended by revolution with many battles fought and lost, and a few won. For surgery at all times has been "up and doing," and "still achieving, still pursuing," it has learned "to labor and to wait." Indeed, it waited for many centuries for the dawn of modern scientific methods. The long night did not begin to break until the middle of the 19th century, when Morton presented to a waiting world the balm of anesthesia. And the bright light of day, bringing in its trail the scientific era of surgery, did not come until Pasteur discovered the cause of wound infection and Lister the possibility of its prevention .

We do well to bear in mind that much of our progress represents a change in method rather than in principle. The basic principles of surgery do not change. The monks of the middle

ages smeared the wounds of their patients with pitch, with more or less beneficial results. Similarly empiric were the methods adopted in subsequent times for keeping wounds clean; today we know the why and wherefor thereof and call it infection, antisepsis and asepsis. Considering the lack of exact knowledge and the utterly primitive facilities at the disposal of the surgeon of old, it is indeed surprising that anyone ever survived an operation. We cannot help feeling that Paré (1510-1590), known as the father of modern surgery, was right when on being asked about the condition of one of his patients, said, "I do the operating, God heals the wound." But by the God-given light in which we are working in this age, a much greater responsibility rests upon the work of human hands, and lack of success cannot so conveniently, as formerly, be shifted, or attributed, to a visitation of God.

Before the advent of the present, the pathological, era of surgery, surgical procedures were mainly concerned with acute injuries, fractures, tumors, the removal of stones from the bladder, ligation of arteries for the cure of aneurisms, etc. Other regions in the body were truly forbidden ground which only a daring few ventured to tread, and some indeed were altogether *terra incognita*. Since the dawn of modern science of surgery, the sign-posts, "No trespassing," have almost disappeared from the path of the aseptic scalpel and there is practically no part of the human organism which in the hands of the adept it cannot reach. Today we remove tumors from the brain, perform delicate operations on the eye, the ear, the nose, the throat and the lungs; we resect the stomach, remove the gall-bladder, extirpate a kidney, suture the liver and the heart, reunite severed nerves and arteries, and, to some extent, restore lost tissues and replace bones by which form is conserved and function reëstablished in deformed and diseased parts of the body.

We owe the removal of the barriers which formerly blocked the path of surgery to the development and the triumphant entry into its realm of the pathology of the living, along with its sister sciences—bacteriology, serology, experemintal physiology, bio- and physico-chemistry, radiology, etc. This does not imply that we are through with the study of anatomy, for now, as always, the road to good surgery leads to and from the dissecting room. The difference is that today we are in a position to compare normal and pathological conditions in the living subject, and are thus gaining a better insight into the nature and cause of the disease and possible prevention.

This subject of the pathology of the living likewise never loses its interest for the surgeon, provided he is not merely an operator, but a student. It has contributed more than anything else to the advance of the medical sciences and the saving of human life. The possibilities opened up by observations on the living subject are as long as the chain of progress from the dark beginnings of surgery down to our own epoch with its almost phenomenal opportunities and achievements. To this study we owe the clarification of many of our ideas concerning such disorders as duodenal and gastric ulcers, disease of the gall-bladder, the bile passages and the pancreas, tuberculosis, and so through the list.

In disease of the gall-bladder, for example, it is possible to trace the steps in the progress of the disorder from its first infection through the activity of micro-organisms, through the stage of acute and chronic inflammation with its attending complications, gall-stone formation and possible precancerous and cancerous change. There is at present considerable propaganda which claims that opening up the abdomen is not the most rational means of establishing the cause of persistent abdominal symptoms; that by means of such devices as the duodenal bucket, disease of the biliary passages can be definitely diagnosed and eventually cured. The experimental work of the late lamented Meltzer, in demonstrating the action of certain agents that are carried into the duodenum and their influence on the sphincter of Oddi have been long familiar to all of us. I believe, also, there are many who will agree with me that these experiments, interesting in themselves, have not been fruitful of much practical good, although there are some enthusiasts who are building card houses on the basis of these observations. No one more than myself would hail with delight the means of curing disease of the gall-bladder without operation; but for the present I believe that is the only means of making the diagnosis with certainty in otherwise obscure cases and of treating them with a great probability of cure. No one more than myself likewise is conscious of the stubbornness of biliary infection

and how long it sometimes takes to obtain a cure by direct drainage; yet we know that our results, once obtained, with the exception of a small percentage, are not only permanent, but brilliant.

One of the startling facts demonstrated by the study of disease in its progress instead of at its end-result on the autopsy table, is the disposition to cancerous degeneration of chronic ulceration of the pyloric antrum. Our increasing knowledge of cellular pathology in neoplastic tissue removed at operation has entirely changed our views with regard to cancerous changes. This is one of the reasons why excision of duodenal and gastric ulcer is a common surgical procedure today. Indeed, what we know of the clinical history of these ulcers we owe to the systematic coördination of clinical manifestations with the pathology seen at the operating table. Examples of the contribution, made by thi smeans of study to our knowledge of surgical diseases of the abdominal viscera alone could be multiplied almost *ad infinitum*. I trust that the mere glimpses I have presented in these rather disjointed remarks of the importance of this ever-fascinating and interesting topic may be of value in stimulating the desire to trace symptoms of certain diseases, especially of the gastro-intestinal tract, to their actual physiological and pathological causes. It is only by this means of proper coördination and rational treatment that we can hope to strike from our register chronic invalidism from diseases that are curable if given the proper study and consideration based on the study of living pathology instead of on disease in its terminal stage.

Original Articles.

BASAL METABOLISM AS AN INDEX OF TREATMENT IN DISEASES OF THE THYROID.*

BY FRANK H. LAHEY, M.D., BOSTON,
AND
SARA MURRAY JORDAN, PH.D., BOSTON.

THE study of metabolism is the investigation of the changes occurring in substances which pass through or become incorporated in the body cells, and of the changes in energy which accompany these processes. In other words, metabolism means chemical changes or changes in the material composition of the body (Stoffwechsel), and dynamic changes or changes in energy (Kraftwechsel).

In 1780, Lavoisier and Laplace formulated and proved the hypothesis that life is a process of oxidation supported by respiration. The experimental proof was the measurement of the heat produced by a guinea pig in an ice chamber, and of the carbon dioxide given off by the same animal when placed for a certain time under a glass jar. Lavoisier's assumption that the heat measured was produced by carbon and hydrogen burned in the lungs and excreted in water and carbon dioxide was later corrected when Liebig, in 1842, and his disciples classified the food-stuffs and showed that heat was produced by the oxidation of protein, carbohydrate, and fat in the cells of the organism.

The second great advance was made in 1851, when the law of conservation of energy, then being applied by Helmholtz to all the realms of physics, was shown by Mayer to be applicable likewise to the production of heat by the human body. It was thus demonstrated that energy and matter have the same behavior in the realm of physiology as in that of inanimate nature. Furthermore, it was found that in addition to heat produced by the animal body, the work of muscle contraction, electricity, and even light, as in the firefly, resulted from the transformation of the potential energy of the intake. Later, in 1857, followed the work of Carl von Voit, who first definitely proved that the nitrogen excretion of urine and feces could be used as a measure of the protein in metabolism. Pettenkofer and Voit demonstrated with their respiration apparatus and their chemical analyses of the excretions that the protein, fat, and carbohydrate of the intake could be measured in the carbon dioxide and nitrogen excretions.

Rubner, in 1885, and subsequent years, established the standard fuel values of 1 gram of protein, 1 gram of fat, and 1 gram of carbohydrate as 4.1, 9.3, 4.1 calories, respectively. These values are in use today. With their aid and with the application of the law of conservation of energy, it became easy to calculate the quantity of heat which under normal conditions would be produced by the oxidation of a certain amount of absorbed food-stuffs in 24 hours. When the unabsorbed food-stuffs in feces and

* We wish to express our appreciation to Dr. Thorne M. Carpenter of the Carnegie Nutrition Laboratory, and Miss Edna Tompkins for advice and assistance in the development of the technique of this test.

urine were analyzed and subtracted from the intake, the calculated heat was found to correspond exactly to the heat actually given off.

It was now clear that the metabolism of man could be measured. Then began the long and productive series of investigations of the metabolism of single substances, of single organs or tissues, and of the animal organism as a whole. In the study of the metabolism of the human organism, the variations dependent upon age, sex, height, weight and composition of the body, the influences of temperature, sleep, and other conditions, the great variations in normal metabolism, under the influence of increased or decreased functional activity and chemical substances—the factors which alter metabolism—have all been carefully investigated. The results of these investigations show very definitely that although metabolism may not yet be a "precisely exact science," yet it is possible to express its rate in figures, and in the words of Robert Mayer, "One figure has more true and permanent value than a precious library full of hypotheses." To be sure, figures have their true and permanent value only when obtained by a method which combines accuracy of measurements with a definition of experimental conditions.

The investigations of the metabolism of man have dealt with quantitative variations of normal processes and of pathological conditions as compared with normal. In the comparison of pathological with normal metabolism, a definition of experimental conditions is indispensable. The definition of experimental conditions in the measurement of metabolism has come to be known as the measurement of "basal metabolism." Basal metabolism was expressed by Rubner as "Grundumsatz" or "Ruheumsatz" and in English has been defined as the condition of minimal of fundamental exchange, the irreducible minimum of exchange of energy which is characteristic for the individual. It is the minimal activity of the body which maintains the functions of respiration, circulation, and secretion. Magnus Levy and Johannson have shown that the variations for one individual are very slight under normal conditions.

In the last twenty-two years it has been recognized that in a test of the rate of basal metabolism, digestion and voluntary movements must be eliminated. To accomplish this the patient is tested twelve to fifteen hours after the ingestion of food and while completely at rest. Digestion is eliminated because it has been found that the presence of food in the alimentary canal stimulated metabolism. The importance of complete muscular repose and relaxation is shown by the fact that very slight voluntary and reflex movements or even the maintenance of an uncomfortable position requiring the tension of certain muscles raise the metabolism ten to twenty per cent. or higher. Complete relaxation of the body of the patient in a comfortable reclining position, the most thorough avoidance of all movements, together with his abstinence from food for at least twelve hours previous to the test, are the necessary conditions for a test of his basal metabolic rate. External influences, such as abnormal light, noise, physical or psychical stimuli, should be avoided, since it has been found that slight deviations from the basal rate are produced by the reflexes (respiratory, cardiac, and muscular) which such stimuli elicit.

Two methods have been developed for measuring metabolism. Direct calorimetry, the direct measurement of the heat evolved by the absorbed food-stuffs, was elaborated and applied to man in the calorimeter of Atwater and Rosa, with further developments by Benedict. An indirect method, a determination of the respiratory exchange, has also been developed by which the measurement of oxygen absorbed or carbon dioxide excreted, or both, can be used to calculate the total heat production.

The basis upon which this method was established is the fact that definite amounts of absorbed oxygen are used and likewise definite amounts of excreted carbon dioxide and of evolved heat are obtained by the oxidation to their end-products of food-stuffs of definite composition. It is therefore clear that a certain volume unit of oxygen or of carbon dioxide corresponds to a certain definite amount of liberated heat according to the substance oxidized. This definite amount of heat is the calorific value of oxygen or of carbon dioxide. A liter of oxygen has a calorific value in the combustion of protein of 4.600 calories, in the combustion of carbohydrates, 5.047 calories, and in the combustion of fat, 4.686 calories. A liter of CO_2 derived from the combustion of protein, carbohydrate, and fat, corresponds respectively to 5.683 calories, 5.047 calories, and 6.694 calories. When the amount of oxygen absorbed or of carbon dioxide excreted by an individual is known for a certain period of time, the cal-

ories produced by him can be computed by using a certain calorific value for oxygen or carbon dioxide, which value depends on the nature of the substances metabolized. (The choice of calorific value will be discussed below.

This indirect or gasometric method of calorimetry has been developed in two ways: (1) for the measurement of excreted carbon dioxide and absorbed oxygen, for longer periods of time (up to 24 hours), as illustrated by the apparatus of Rubner, Pettenkofer and Voit, Tigerstedt-Sonden and Jacquet; (2) for the measurement of gas exchange in the lungs, *i.e.*, the absorption of oxygen and exhalation of carbon dioxide for brief periods of time (ten minutes to one hour). The latter is the older method of the French and English scientists, elaborated by Speck (1892), and developed by Zuntz-Geppert (1905), and by Benedict in his "Universal Respiration Apparatus" (1912). These apparatus are of two types, the closed circuit and the air current types. As an example of the latter, the widely used apparatus of Tissot (1898), (1904), may be mentioned, in which a spirometer is used to record graphically the expired volume of air from which a sample is later analyzed.

An example of the closed circuit type of apparatus, which has the advantage of requiring no gas analysis for the computation of results, is that used in this present series of tests, the Benedict Portable Respiration Apparatus, devised by Benedict in 1918, and in use in the laboratory at the New England Deaconness Hospital since October, 1919.

The apparatus is described by Benedict in the BOSTON MEDICAL AND SURGICAL JOURNAL, May 16, 1918. The essential feature is the connection between the lungs of the patient and a spirometer, filled with a mixture of room air and oxygen. This connection is made by means of a rubber mouthpiece which connects the mouth of the patient with pipes leading to and from the spirometer. Nasal breathing is shut off by the use of a nose clip. A closed circuit is thus established which includes the lungs of the patient, the spirometer which rides up and down on water with each respiration, and two jars of sodium hydroxide which are placed within the circuit to absorb the carbon dioxide as it emanates from the lungs. An electric motor is used to blow the air around the circuit.

The absorption of oxygen by the lungs of the patient is measured and can be directly read

CHART I.

RATE OF BASAL METABOLISM.

Name—Miss J. A. F. *Date*—Sept. 29, 1920

Date of birth—July 5, 1898 (22 yrs.)
Body weight—132.5 lbs. (61.33 kilo.)
Height—162.2 cm.
Surface area—1.65 sq. M.
Buccal Temp.—98.2°

No. of period—1 { Period begun—7.22 A.M.
Duration of period—9 min. 2⅖ sec. (9.043 min.)

Barometer Reading		*Temp. of Spirometer*	
Log. 769	2.88593	Begin	27.5° C.
Log. 760	2.88081	End	28.0° C.
	0.00512	Av.	27.75° C.

Pulse	*Respiration*	*Temp. Reduced to Standard*
(96)	(18)	
86	12	Log. 273 .. 2.43616
96	12.5	Log. 300 7—2.47813
100	14	
98	15	9.95803—10
96		

Readings of Height of Spirometer

Beg.	7170	
2 min.	6600	570
4 min.	6000	600
6 min.	5400	600
8 min.	4820	580
9 min. 2⅖ sec.	4470	350
	2700 — 2700 =	Diff. in readings of height of spirometer for period.

Log. diff. height spir.	3.43136		
Log. temp. standardized	9.95803—10		
Log. bar. standardized	0.00512		
Log. vol. oxygen absorbed ...	3.39451		
Log. duration of period	0.95631		
Log. vol. O_2 absorbed per min.	2.43820	=	274.3 c. c.
Correction for total vol. air ..			1.0 c. c.
			275.3 c. c.

CHART I.—A post-operative observation. Pulse and respiration readings in parentheses were made just before the period was begun. The others were made at two-minute intervals during the test. For control during the test, the differences in the two-minute readings of the height of the spirometer are quickly calculated and compared. The patient is allowed to breathe into and out of the apparatus for 1.5 minutes before the first reading is made. Spirometer readings are made at the end of expirations. A correction of 2 c.c. for each degree C. rise in temperature is made for reduction of total volume of air to 0 degree C. and 760 mm. pressure.

Readings are made of at least two periods of about ten minutes each. If the results are found to vary too greatly from each other, further readings are made.

on the scale of the spirometer (Chart I). The excretion of carbon dioxide can likewise be determined by weighing, before and after the test, the jars of sodium hydroxide which absorb it. The latter method is, however, not as desirable as the former, inasmuch as the carbon dioxide excretion has been found to be more affected by abnormal respiration than the oxygen absorption, and since, furthermore, the calorific values of carbon dioxide deviate more from one another than do those of oxygen.

As the amount of oxygen absorbed by the patient corresponds to a definite amount of liberated heat, according to the substance oxidized, the calorific value of oxygen is the next consideration. This is dependent upon the relationship of carbon dioxide to oxygen, or the respiratory quotient, which varies according to the nature of the material metabolized. From a study of respiratory quotients by Benedict and others, 0.82 has been found to be an average. The calorific value of oxygen corresponding to this average respiratory quotient is 4.825, as determined by Zuntz and Schumburg. By multiplication of the amount of oxygen absorbed per minute (measured by the spirometer and reduced to a standard temperature and barometric pressure), by the calorific value 4.825 the number of calories produced per minute is obtained. It is now necessary to compare this actual heat production of the individual with a standard of norm (Chart II). Here we are confronted with a situation inevitable in a science which is not "precisely exact." In the words of Dr. Benedict, "There is no inflexible standard for normal metabolism for any given age, weight, height, and sex, from which all normal individuals never vary." The most widely used method for computing normal metabolism has been, in the past, the calculation based on the "body surface law," the underlying conception of which is that the amount of heat produced by the body is proportional to the surface area rather than to the weight. In 1883, Rubner published the results of experiments which seemed to demonstrate the existence of an "emission constant," or constant heat production per unit of body surface. The conception had been held for several decades previous to Rubner's work, and Richet published experimental data in 1889, but the law is commonly known as Rubner's Law. The physiological evidence for the validity of this law has been discussed by Benedict in 1919, and shown to be inconclusive. Nevertheless, this law has been in common use as the basis of the measurement of normal metabolism.

The measurement of the surface area of man has been accomplished in various ways, such as the covering of the body with paper and the measurement of the area of the paper or a protographic method with the measurement of the prints by means of a planimeter. Formulae were made from such measurements by which surface area can be computed from the height and weight of an individual. The linear formula and the height-weight chart of Du Bois and Du Bois, published in 1915, are widely used today.

However, in 1919, Harris and Benedict published the results of their biometric studies of the basal metabolism of man, in which they showed from a statistical analysis of the metabolism of over two hundred normal individuals, together with the correlation of the factors for their body weight, stature, age, and sex, that the proximate factors determining the metabolism are the body weight, height, age, and sex, all of which have an independent influence upon metabolism. From their analysis of these factors they have established new formulae, and from these formulae have made multiple prediction tables for the basal metabolism of the two sexes between the ages of twenty-one and seventy. These tables have been used in this series of tests, and it may be stated that in general the values thus determined conform with the clinical aspect of the cases. Comparisons

CHART II.

SHOWING CALCULATION OF PERCENTAGE VARIATION FROM NORMAL, BASED ON OBSERVATION SHOWN IN PLATE I.

Miss J. A. F.

Actual Basal Metabolism

Vol. oxygen absorbed per min. { Period 1....275.3 c.c. / Period 2....275.1 c.c.

Average .....275.2 c.c.

$\frac{275.2 \text{ c.c.}}{1000 \text{ c.c. (1 litre)}} \times 4.825$ (cal. val. of 1 litre O_2) =

1.32784 calories per minute

1.32784 x 1440 (minutes per day) = 1912.0896 calories per day

Normal Basal Metabolism

1. Harris and Benedict Method (Prediction Tables)

Height and age 197

Weight 1241

1438 calories per day

2. Du Bois Height Weight Chart and Linear Formulas

Surface area ..... 1.65 sq. M.

1.65 x 37 { (Av. calories per hr. per sq. M. of body surface of women, 20 to 29 years) } = 61.05 calories per hr.

61.05 x 24 (hrs. per day) = 1465.20 calories per day

Comparison of Normal with Actual Basal Metabolism

(To obtain percentage variation from normal)

(H. & B.) 1438 : 1912.0896 = 100 : X

X = 132

Percentage variation from normal +32

(Du B.) 1465 : 1912.0896 = 100 : X

X = 130

Percentage variation from normal +30

have frequently been made with computations based on the DuBois height-weight chart and linear formula and slight deviations have been noted (Chart III).

CHART III.

SHOWING VARIATIONS IN ESTIMATED RATE OF BASAL METABOLISM BY COMPARISON WITH HARRIS AND BENEDICT AND WITH DUBOIS NORMALS. (10 CASES.)

NAME	AGE YEARS	HEIGHT IN CM.	WEIGHT IN KILO.	HARRIS AND BENEDICT METHOD	DUBOIS METHOD
Mr. B.	26	166	54.4	+59	+55
Mrs. A.	28	161	45.4	+66	+63
Miss I.	20	151	46.1	—26	—23
Mrs. B.	51	156.5	71.1	—7	—11
Mrs. F.	42	174.5	61.9	—14	—22
Mrs. P.	27	162	46.5	+36	+33
Miss F.	22	162	61.3	+32	+30
Mrs. P.	43	158.5	40.9	+44	+38
Miss F.	25	153	54.7	—3	—4
Mrs. P.	42	153	35.6	+28	+28

Therefore, although it can not yet be stated absolutely that an individual with a basal metabolic rate within ten points of the standard or norm (as determined by either of the above methods outlined), is actually within *his* normal metabolic limits, yet it can be assumed that the deviation from his norm is not great. Until further statistical data are collected, the greater value of the determination of a basal metabolic rate must remain intra-individual, that is, in the comparison of the basal metabolic rates of the same individual under different circumstances, as, for example, under surgical treatment as in this series.

In addition to the possible error arising from comparison of the actual basal metabolic rate with a norm which is relatively indefinite, another source of error lies in the technique of the measurement. The experimental conditions have been described above. To avoid gross errors, these conditions must be strictly observed. Furthermore, the condition of the apparatus itself offers possibilities of error. It must be carefully tested for leakage before each test and the soda lime must be renewed often enough so that it remains an efficient absorber of carbon dioxide. Otherwise, the patient suffers great discomfort from the absorption of carbon dioxide and its presence in the spirometer interferes with the measurement of oxygen. Careful and accurate readings and computations are, of course, also necessary. In the experience derived from this series of tests, it has been found that the chief sources of error in technique arise from two defects in the condition of the apparatus, namely, leakage (usually at the joints, due to defective washers), and non-absorption of carbon dioxide by the soda lime. Either condition produces so considerable a deviation from the correct value as to make the results worthless.

In this series, 304 tests have been made upon 135 individuals, 25 of whom were normal controls.* The other 110 individuals were patients suffering from diseases of the thyroid, namely, simple adenomata, adenomata with secondary hyperthyroidism, primary hyperthyroidism, or hypothyroidism.

* The data discussed in this paper are those gathered up to October 1, 1920.

Basal metabolism as a diagnostic measure, although subject to error in estimation and interpretation, is probably more constantly reliable than many of our other laboratory measures, both as to its qualitative and quantitative value. It is of the greatest value in hyperthyroidism, since that disease is so distinctive in its course and symptoms, and so characteristically a disease of hypermetabolism.

From our experience in metabolism studies in connection with thyroid disease, involving 304 metabolism estimations on 135 patients, it does not seem that active hyperthyroidism exists without an increase in the metabolic rate.

Such study is of great value in separating those cases of neurasthenia and associated neurological and cardiac lesions, such as effort syndrome, which often present signs closely simulating those of hyperthyroidism, in that these conditions show metabolism estimations within normal limits.

It is of value, further, in determining the presence of secondary hyperthyroidism in those cases of adenoma of the thyroid in which the symptoms of hyperthyroidism are not markedly evident. It used to be a source of considerable surprise to us in the beginning of our thyroid experience, before we were accustomed to employ basal metabolism estimation, to find that many patients upon whom we operated merely to remove an adenoma which was either causing pressure or was unsightly, came back within a few months, having gained in weight, and stating that they felt a marked change in their general health. This we now know to have been an improvement from secondary hyperthyroidism which, when well marked, is—with the exception of exophthalmos—similar in every way to primary hyperthyroidism, and no more diffi-

cult to demonstrate. Its presence, however, in mild or pre-states of clinical hyperthyroidism is not easily demonstrated clinically, but even slight degrees of secondary hyperthyroidism are demonstrable by increases in the rate of metabolic activity, examples of which are shown below as pre-and post-operative metabolism determinations (Chart IV).

CHART IV.

SHOWING PRE- AND POST-OPERATIVE BASAL METABOLISM ESTIMATION IN A FEW CASES OF SECONDARY HYPERTHYROIDISM.

NAME	AGE	DATE	PULSE RATE	BAS. MET. RATE	OPERATION
Mrs. B.	25 yrs.	Mar. 18, 1920	118	+55	Excision of adenoma
		Mar. 19, 1920			
		Mar. 30, 1920	108	+32	
		Apr. 27, 1920	86	+12	
Mrs. B.	34 yrs.	Mar. 11, 1920	96	+36	Excision of adenoma
		Mar. 12, 1920			
		Mar. 24, 1920	62	+15	
Miss C.	26 yrs.	Jan. 9, 1920	90	+20	Excision of adenoma
		Jan. 10, 1920			
		Jan. 16, 1920	74	+12	
Mrs. O.	27 yrs.	Mar. 8, 1920	112	+33	Excision of adenoma
		Mar. 10, 1920			
		Mar. 19, 1920	70	+11	
		May 17, 1920	68	+4	
Mrs. Z.	58 yrs.	May 11, 1920	110	+39	Excision of adenoma
		May 12, 1920			
		May 21, 1920	86	−8	

Metabolism estimation correlated with the clinical examination is also of marked value in indicating the ability of a patient to withstand surgical procedures of various magnitudes; for example, in deciding whether the ligation of a single thyroid pole should be employed, whether it would be wise to ligate both thyroid poles, or whether it would be safe to employ partial thyroidectomy as a primary procedure. It is of further value in deciding when to follow the preliminary pole ligation with the partial thyroidectomy. Up to recently it has been our custom to ligate the superior thyroid poles and then send the patient home for eight weeks, at the end of which time the patient returned to the hospital for the complete operation. As the result of our metabolism investigations, we have been finding that in some cases by waiting eight weeks, we have passed the period of maximum improvement, as indicated by a rise in the metabolism rate from the rate representing the maximum drop after pole ligation. We have, therefore, come to the conclusion that during the intervals between pole ligation and partial thyroidectomy, metabolism estimations should be made every two weeks, and partial thyroidectomy employed as soon as the maximum gain indicated by a drop in metabolism and pulse rate is made.

Since we do not feel that metabolism rate alone in hyperthyroidism is a true criterion of the seriousness of a patient's condition, and, considered alone, is unreliable as a guide to the extent of the surgical procedure to be employed, it is at once evident that it is impossible to make any fixed statement concerning the percentage figures in basal metabolism at which it is safe to perform partial thyroidectomy as a primary procedure. Neither is it possible to state at what level of basal metabolic rate it is wise to ligate but one pole, nor is it further possible from basal metabolism estimations alone to weed out the cases too seriously afflicted to be submitted to any surgical procedure, and to be relegated to the less satisfactory and more prolonged methods of treatment not surgical in nature, such as rest and x-ray treatment.

Yet with the previous view clear in our minds, we are willing to make certain approximate groups which we believe, from our experience, work out satisfactorily in a majority of instances.

Partial thyroidectomy may be employed as a primary measure, unpreceded by ligations, in most of the cases showing increases in metabolic rates of not over +35. In all such cases, however, careful consideration must be given to the apparent degree of toxicity of the disease, as evidenced by the clinical signs, and this consideration must not be biased by the degree of increase in basal metabolic rate, however moderate it may be. Careful consideration must also be given resulting or associated lesions, such as myocarditis and renal lesions.

Fine judgment, based largely upon an extensive experience as to how similar cases have acted under various surgical procedures, is necessary in deciding between pole ligation and primary partial thyroidectomy in the cases of hyperthyroidism showing increases in basal

metabolic rates of between +35 and +50. Decision as to the type of operation should be left until the patient is on the operating table and anaesthetized, and should then be made only when based upon the known basal metabolic rate, the pulse rate, its character and rhythm, and with a consideration of the possible degree of post-operative thyroid reaction. It is our enstom to make this decision with the coöperation of our anesthetist, Dr. Lincoln F. Sise, who in 500 operated thyroid cases has had the same opportunity as we have had to observe their condition before, during, and after various surgical procedures. We have but one fixed rule in deciding as to the wisdom of primary partial thyroidectomy. When doubt exists even in the slightest degree as to the safety of primary partial thyroidectomy, we ligate. We do not permit ourselves to be influenced by the possible disappointment of the patient or her friends in that a second and possibly a third and a fourth operation may be needed before permanent relief is accomplished. We are unbiased by the consideration that the patient may never return for the final operation. Once or twice we have been biased by the fact that a patient had came from a long distance, such as the Middle-West or the provinces of Canada, and was anxions to be saved the cost and discomfort of either a trip home and return within two months, or a stay of perhaps two months away from her home. Twice have we permitted these considerations to sway us in favor of primary partial thyroidectomy. In both instances severe reactions occurred from which the patients, to our relief, recovered; but no longer are we influenced in the least by any of the foregoing considerations. Where life lies in the balance, judgment as to the procedure, in our opinion, must be based entirely upon facts which have to do only with the limiting of risk to the individual.

Most patients showing basal metabolic rates above +50 are safest when submitted to primary ligation of one or both of the superior thyroid poles. It is our custom in these cases to reserve decision, as previously stated, until the patient is anaesthetized (scopolamine, morphia, novocaine + gas oxygen), making use of the same measures as stated above. The decision is then made whether the patient shall be sent back to bed to observe the degree of reaction to the anaesthesia and visit to the operating room, or whether ligation of the poles shall be proceeded with. If the latter procedure is decided upon, one superior pole is rapidly ligated and decision as to whether ligation of the other superior pole is to be carried out is based upon the immediate condition of the patient and how she has endured the ligation of a single pole.

In patients showing metabolic rates of from +75 up, extreme caution must be exercised in the extent of the surgical treatment made use of. In the doubtful cases, a visit to the operating room and gas oxygen anaesthesia with no operative procedure gives valuable evidence as to the resulting degree of reaction. In the less doubtful cases, the ligation of a single pole with a wait of ten days to two weeks makes the ligation of the second pole very much more safe for the individual (Chart V).

CHART V.

SHOWING PROGRESSIVE DROP AS RESULT OF A SINGLE POLE LIGATION.

NAME	AGE	DATE	PULSE RATE	BAS. MET. RATE	OPERATION
Mrs. F.	53 yrs.	Jan. 24, 1920	130	+99	
		Jan. 29, 1920			Double sup. pole ligation
		Mar. 31, 1920	120	+74	
		Apr. 1, 1920			Single inf. pole ligation.
		Apr. 14, 1920	104	+36	
		Apr. 15, 1920			Partial thyroidectomy
		Apr. 25, 1920	106	+4	
Miss D.	20 yrs.	May 27, 1920	146	+60	
		May 28, 1920			Single sup. pole ligation.
		June 3, 1920	136	+38	
		June 4, 1920			Sec. single sup. pole ligation
		July 29, 1920	134	+45	
		July 30, 1920			Partial thyroidectomy
		Aug. 25, 1920	112	+29	

In case (Miss D.), note increase in metabolic rate between observations of June 12th and July 29th from +10 to +45, a possible example of a delay beyond the optimum point following ligation.

Chart VI.

Chart Showing Metabolic Rate During Course of Surgical Treatment in Ten Cases of Primary Hyperthyroidism.

Name	Age Yrs.	Date	Pulse Rate	Body Weight Lbs.	Bas. Met. Rate	Operation
Mrs. A.	34	May 14, 1920	106	123.2	+55	
		June 22, 1920	114	116.6	+61	
		June 23, 1920				Double sup. pole ligation
		June 28, 1920	106	112.2	+57	
		Aug. 3, 1920	104	116.9	+21	
		Aug. 11, 1920				Partial thyroidectomy
		Aug. 19, 1920	74	114.7	+1	
Mrs. A.	53	Dec. 15, 1919	128	98	+77	
		Dec. 16, 1919				Double sup. pole ligation
		Mar. 18, 1920	122	90.5	+58	
		Mar. 19, 1920				Partial thyroidectomy
		Mar. 28, 1920	92	87.5	+19	
Mr. A.	23	Apr. 15, 1920	120	117.2	+81	
		Apr. 16, 1920				Double sup. pole ligation
		June 16, 1920	108	120.2	+60	
		June 18, 1920				Partial thyroidectomy
		June 25, 1920	80	113.9	+34	
		Sept. 4, 1920	54	132.9	—16	
Miss B.	25	May 19, 1920	118	109.7	+62	
		May 20, 1920				Double sup. pole ligation
		May 27, 1920	112	104.6	+30	
		July 14, 1920	104	103.3	+34	
		July 15, 1920				Partial thyroidectomy
		July 24, 1920	94	97.6	+8	
Miss D.	28	Jan. 8, 1920	120	125.1	+93	
		Jan. 9, 1920				Double sup. pole ligation
		Jan. 15, 1920	106	119	+68	
		Mar. 9, 1920	98	125	+58	
		Mar. 10, 1920				Partial thyroidectomy
		Mar. 21, 1920	74	128	+15	
		July 15, 1920	74	140.2	+17	
Mrs. F.	28	Oct. 15, 1919	130	119.5	+63	
		Oct. 26, 1919				Double sup. pole ligation
		Nov. 2, 1919	124	110.7	+72	
		Jan. 22, 1920	116	111.2	+55	
		Jan. 23, 1920				Partial thyroidectomy
		Feb. 1, 1920	86	102.7	+24	
Mrs. H.	58	Feb. 4, 1920	90	145	+37	
		Feb. 5, 1920				Double sup. pole ligation
		Feb. 20, 1920	94	137	+27	
		Apr. 10, 1920	82	146.6	+11	
		Apr. 11, 1920				Partial thyroidectomy
		Apr. 24, 1920	76	140	—5	
Miss H.	44	Dec. 11, 1919	124	103	+70	
		Jan. 3, 1920				Double sup. pole ligation
		Jan. 11, 1920	120	93.7	+82	
		Mar. 2, 1920	98	113.7	+21	
		Mar. 3, 1920				Partial thyroidectomy
		Mar. 14, 1920	116	100.7	+30	
		July 9, 1920	94	108.5	+23	
Miss P.	22	Dec. 29, 1919	130	84.3	+72	
		Dec. 30, 1919				Double sup. pole ligation
		Jan. 12, 1920	96	77.2	+100*	
		Mar. 5, 1920	106	91.5	+23	
		Mar. 6, 1920				Partial thyroidectomy
		Mar. 15, 1920	96	81.5	+29	
		Apr. 30, 1920	84	96.2	—5	
Mrs. S.	44	July 2, 1920				Double sup. pole ligation
		July 10, 1920	120	120.2	+79	
		Aug. 11, 1920	106	134	+51	
		Aug. 27, 1920	114	137.5	+53	
		Aug. 28 1920				Partial thyroidectomy
		Sept. 3, 1920	94	129.7	Normal	

Note.—It is to be noted that in some cases the immediate effect of ligation is a temporary rise in the metabolic rate.

* Fever

It is our opinion that there is hardly a surgical disease in which caution is as necessary as in the cases of marked thyrotoxicosis. As they approach the higher levels of toxicity as indicated by clinical signs and increases in metabolic rates to high level, each case should be individualized and no longer looked upon as being a member of a group. After such cases have reached the more intense stages of thyroid intoxication, they no longer conduct themselves by rule. This statement is more or less true of hyperthyroidism as a whole, but is particularly characteristic of the intensely toxic cases. After an extensive experience with thyroid cases, we feel able to prophesy with relative certainty the course and reaction of mild and moderate degrees of thyroid intoxication to various surgical procedures, but have by no means the same degree of assurance regarding the intensely toxic cases. It is because of this unknown capacity for unfavorable reaction on the part of severely toxic cases that extremely cautious advance should be made. First, the degree of reaction to the slight discomfort and apprehension which goes with the estimation of basal metabolism, next the reaction to a visit to the operating room and short gas oxygen anaesthesia, and finally based upon this data, the ligation of a single thyroid pole. Having advanced so far, we have accumulated sufficient data upon which to base with reasonable certainty a decision as to the probable outcome with further surgical procedures.

We feel that there will be a tendency to accept basal metabolic rates as true percentage estimates of thyroid hyperfunction, which in turn afford a basis for a true prediction of reactions to surgical proceedures; and we must emphasize our conviction that such a condition of affairs would be extremely dangerous. With the first view, however, we feel some sympathy. That thyroid toxicity, if we may for practical purposes consider the disease in terms of toxic thyroid products, is represented by increased basal metabolic rates there can be little question. One has but to observe the charts shown here, and note the metabolic rate before operation,—the progressive drop in the metabolic rate and pulse rate after ligation and again after partial thyroidectomy, together with the progressive clinical improvement in the patient, going hand in hand in most instances with the drop in metabolic rate,—to be convinced that metabolic rate quite accurately represents degree of toxicity (Chart VI).

Cases have arisen, however, in which there has been marked clinical improvement and gain in weight without proportionate drop in metabolism (Chart VII).

CHART VII.

SHOWING TWO CASES IN WHICH, AFTER LIGATION, CLINICAL IMPROVEMENT AND GAIN IN WEIGHT OCCURRED WITHOUT PROPORTIONATE DROP IN METABOLISM.

NAME	AGE YRS.	DATE	PULSE RATE	BODY WT. LBS.	BAS. MET. RATE	OPERATION
Mrs. C.	40	Mar. 6, 1920	116	107.5	+45	
		Mar. 10, 1920				Sup. pole ligation
		Mar. 16, 1920	108	104.2	+53	
		May 8, 1920	102	115.1	+52	
		May 9, 1920				Partial thyroidectomy
		May 10, 1920	94	108.5	+37	
		July 17, 1920	72	120.7	+15	
		Apr. 23, 1920				Single sup. pole ligation
Miss M.	35	May 6, 1920	130	109.7	+48	
		May 7, 1920				Second sup. pole ligation
		May 12, 1920	124	109.6	+31	
		July 6, 1920	118	118.5	+46	
		July 7, 1920				Partial thyroidectomy
		July 17, 1920	96	115.7	+14	

Regarding the second view—that basal metabolism represents true terms of toxicity which in turn may be used to estimate ability to withstand surgical procedures, we feel very strongly that the level of basal metabolic rate alone is quite undependable, and in fact often misleading as a means of prognosticating the seriousness of the patient's state and her ability to endure any surgical procedure. We have had surgical experiences with patients showing metabolic rates at both extremes, in which the course and results of the operative procedure have been entirely out of accord with what might have been expected from the metabolic rate.

We have, for example, operated upon patients with a basal metabolic rate of +26 with a hyperthyroidism of long standing, and seen an extremely well-marked operative and post-opera-

tive reaction. On the other hand, we have operated upon patients with metabolic rates around +60 or over, but with a hyperthyroidism of short duration, who have shown much less operative and post-operative reaction than have the patients with a rate of +26 as spoken of above.

It is our feeling that one must consider—in estimating the operative risk—not only the degree of toxicity, but also the cardiac reserve as estimated by the rate, rhythm, and character of the pulse wave, by the length of time the disease has been present, and by the history and signs of past or present indications of unsettled compensation, and the destructive effect of the disease on other structures, as indicated by varying degrees of nervousness, tremor, and loss of weight.

Errors of estimation and the avoidance of their causes have been discussed in the first portion of this paper. Given an accurate estimation carefully checked against those influences (principally food and exertion) which in themselves produce increased rates, errors then occur for practical purposes but rarely in the interpretation of these findings. Care must be taken, for example, in thyroid cases, to ascertain the presence of complicating conditions which seriously affect the metabolic rate. For example, diabetes, in our experience, has not been a rare complication of hyperthyroidism (or *vice versa*, if you choose), and in such a condition care must be taken to ascertain to what diet and to what treatment the patient has been subjected.

Chart VIII.

Showing Effect upon Basal Metabolism Rate of Injection of 8 Minims of Epinephrin (1-1000 Sol.).

Mr. H., Age 53 Yrs.	Bas. Met. Rate	Pulse Rate	Blood Press.
Before injection	+76	102	130/88
Minutes after Injection			
5	+77	108	132/90
11		112	148/88
15		107	142/94
20		120	166/86
25		120	168/86
30	+89	119	176/102
35		120	160/90
40		124	146/96
47		124	138/94
55		124	134/96
60	+78	124	134/96
65		116	134/94
70		116	132/92

Care must also be taken to ascertain possible renal lesions, marked examples of which are said to give increased metabolic rates.

Seemingly simple infections giving but low degrees of temperature reaction must also be eliminated, as fever very promptly results in a marked increase in metabolic rate.

In thyroid cases, basal metabolism estimation should not be undertaken at a time too close to the period following the performance of the Goetsch or epinephrin test, as injections of epinephrin have resulted, in our experience, in well-marked rises in metabolic rates (Chart VIII).

Apprehension, anxiety, and excitation play a considerable part in the production of unreliable data in metabolism estimation, and where even the possibility of such influence occurs, the test should be repeated daily until an average is struck or the patient eliminated as a subject from whom a reliable result may not be obtained.

Chart IX.

Showing Average Rate of Basal Metabolism in Secondary and in Primary Hyperthyroidism.

Adenoma with secondary hyperthyroidism (25 cases)	
Before excision of adenoma	+40.44
After excision of adenoma (av. 9.5 days)	+22.61
Primary hyperthyroidism (25 cases). Preliminary ligation of poles	
Before ligation	+62.00
After ligation (av. 7.5 days)	+52.19
Before partial thyroidectomy (av. 61 days after ligation)	+43.84
After partial thyroidectomy (av. 9.1 days)	+19.47

Out-patients presenting themselves for estimation upon the morning of the test may or may not have observed the essential necessity of having abstained from nourishment or stimulants for fourteen hours previous to the test, so that for accurate results it should be insisted that the patient spend the night before the test in the hospital.

Associated lesions of the pituitary, wasting disease, prolonged dieting, particularly for diabetes, are thought to be some of the possible causes of abnormally low readings.

When all the above-mentioned factors can be eliminated and the basal metabolic rate can be regarded as a true index of thyroid activity, it may be said in conclusion, that, in our opin-

ion in a clinic for diseases of the thyroid, a test of the rate of basal metabolism is an invaluable factor in the data necessary for diagnosis; secondly, that it aids greatly in prognosis; thirdly, that it probably has its greatest value as an index of the results of treatment in any given case.

Medical Progress

REPORT ON RECENT PROGRESS IN DERMATOLOGY.

By John T. Bowen, M.D., Boston.

POST-VACCINAL RINGWORM (TRICHOPHYTOSIS?)

Gougerot[1] states that the examination of thousands who were vaccinated during the war has shown that ringworm, although rare, was the most frequent complication of vaccination, and the most important in practice. Other complications were impetigo and erythema, very rarely septic or other affections. The subjects were of all ages, soldiers, children, workers in powder factories, and the affection usually began at the end of the vaccinal process when the crust was about to fall, from the fifteenth to the twentieth day. The development was usually slow, in one case, a year; sometimes, however, rapid, one patient having, eight weeks after the vaccination, the arm almost entirely covered and the chest spotted with foci of autoinoculation. There is nothing peculiar about the clinical aspects, the forms varying according to the variety of the species inoculated. The process seems to go on indefinitely if not treated. Gougerot declares that negative bacteriological results should not deter one from a clear clinical diagnosis. The treatment, the same as that of trichophytosis of the smooth skin, although simple, is entered into with much elaboration and detail by the writer.

Etiologically, it is to be supposed that the vaccine was uncontaminated and that the skin abrasion offered the gate of entrance to the fungus contained in the clothing or existing on the skin of certain persons. In other instances, family or atelier epidemics seem to prove a contamination of the virus.

With regard to prophylaxis, it seems that contamination of the vaccine could be eliminated by careful examination of the skin of the heifers before inoculation, both clinically and microscopically. Gougerot thinks the subject of importance in that these complications are very often misinterpreted. He disclaims any desire to criticise the excellent French vaccination plants, or to raise this slight complication as an argument against vaccination.

THE PARASITIC NATURE OF DYSIDROSIS.[2]

Of late the impression has been steadily gaining ground that at least many of the cases described under the heading of dysidrosis, by Tilbury Fox, and of pompholix by Hutchinson, are really a form of epidermophytosis of the extremities. Darier noticed during the war that whenever a soldier appeared with a vesiculobullous or squamous eruption limited to the hands and feet, three diagnoses were usually considered,—eczema, occupational dermatitis, or dysidrosis—only exceptionally was a parasitic mycotic affection thought of. Diathetic and seborrhoeic eczema are seldom, if ever, localized on the hands and feet; whereas the professional or artificial forms of dermatitis have their seat on the hands, less often on the feet, and are recognized by the character of the initial lesions and by inquiry about the occupation. Tilbury Fox's dysidrosis and Hutchinson's cheiropompholix, or simple pompholix, is a vesiculobullous dermatitis, later on squamous in type, essentially limited to the hands and feet, and comes on in recurring attacks, especially in spring and summer; is symmetrical, is accompanied by burning or tingling, and often appears after profuse perspiration, a nervous shock, or violent exercise, and is said to be most common in overworked, dyspeptic, neurasthenic subjects.

It has been found in the last fifteen years that various epidermycoses can produce quite the same appearance as dysidrosis and eczema of the extremities. A palmar and plantar trichophytosis was first described in 1892 by Djelaleddin Mouktar, and differentiated from dysidrosis. Then came the work of Sabouraud from 1907-1910 on epidermophytosis, an affection due to a parasite closely allied to the trichophyton, the epidermophyton inguinale, which does not attack the hairs. When situated in the groin and on the upper part of the thighs, it causes the form known as Hebra's eczema marginatum, showing very red, round or polycyclic patches, rapidly extending, with sharply bounded edges, and sometimes vesicular. Occasionally this form is seen in the arm-

pits, and other parts of the trunk. It is very common in the interdigital space between the toes, showing a macerated epidermis, covering a red base and sometimes accompanied by vesicles. It may reach the instep, and may continue indefinitely. On the hands it is rarer, exhibiting irregular vesicles on the lateral surfaces of the fingers, without forming circles, and can be surely identified only by a microscopical examination which shows in the scales and vesicles the characteristic mycelium. Sabouraud pointed out that the etiological conditions to which dysidrosis is attributed were very common, whereas numerous cases of this form were not recorded. He found among soldiers attacked by vesiculo-bullous or squamous eruptions on the hands and feet: (1) A huge majority of secondary eczema from scabies, pediculosis, boils, wounds, etc. (2) A considerable number of cases of occupational dermatitis. (3) A very few cases of palmar trichophytosis. (4) A moderate number of cases which best fitted in the category of dysidrosis. The types of these were: (a) Vesicular and bullous type, clear vesicles scattered or vaguely grouped on the palmar and lateral surfaces of the fingers, on the palm, and sometimes on the back of the hands; lenticular and bullous lesions, often purulent, on the forepart or whole sole of the foot. (b) Vesiculo-squamous type, in which the lesions were made up of erythemato-squamous blotches, scattered or confluent, with crumbling edges. (c) Squamous type, dry desquamations, irregular with wavy edges, only exceptionally circinate or polycyclic. This is complicated constantly with intertrigo of the toes.

In order to determine how many of these cases are epidermic mycoses, Darier examined the roofs of the vesicles and the scales microscopically. He found the epidermophyton in such a large proportion of the lesions examined that he is forced to the conclusion that dysidrosis does not exist as a distinct cutaneous disease, and that the clinical picture so well portrayed by Fox and Hutchinson is that of epidermophytosis of the extremities. His method was to treat the shreds of epidermis with alcohol-ether to remove the fat, then separate by needles in acetic acid, and finally clear on the slide in 40% caustic potash with gentle heat. He admits, however, that it will be necessary for further investigations to confirm this conclusion and to determine the exact nature of the parasite and whether one or several species are concerned. Furthermore, the clinical facts support the view that dysidrosis is a mycotic disease, inasmuch as it is said that the attacks of dysidrosis usually last from ten to twenty days and often recur in the same subject, and that this is true of epidermophytosis of the extremities; but it must be added that one also sees continuous attacks prolonged for six months. The two affections are much alike, also, in topographical distribution: attacking hands, feet, or all four extremities symmetrically at the same time.

EXFOLIATIVE DERMATITIS DUE TO ARSPHENAMINE.

Latham,[3] at Columbia, S. C., reports that there had occurred during the preceding five months four cases of this affection of which he describes the most typical—a fatal one which followed the intravenous injection of 0.8 gm. of arsphenamine. A man of 27 had a positive Wassermann, but exhibited no signs or symptoms of syphilis and denied a venereal history. He received an injection of 0.4 gm. of arsphenamine intravenously, and a second dose of the same amount one week later. Ten days after the second dose he was admitted to the medical ward of the base hospital with an intense erythema and beginning infiltration of the skin of the face and neck, with superficial desquamation and pigmentation, together with some conjunctivitis. At this time there was no evidence of polyneuritis. Arsenic was found in the urine, but apparently had excited no renal irritation. The erythema and exfoliation gradually became generalized and a moderate fever appeared. There was some fissuring and exudation. Four weeks after the first dose of arsphenamine, the only constitutional symptoms were a moderate fever and a pronounced leucocytosis. As the exfoliation progressed, toxic symptoms appeared, the pulse and fever rose, and a laryngobronchial catarrh asserted itself. Mental symptoms appeared coincidently with some sensory disturbance and stupor; coma and light convulsions were present. There was increasing soreness of the mouth, so that toward the end he had to be fed by a tube. There was evidence of severe trophic disturbance of the blood vessels causing extraversation of the blood from the slightest wound. Thirty-four days after the first dose of arsphenamine, he was in a state of coma, with a definite bronchopneumonia. He died on the thirty-eighth day.

The urine showed traces of arsenic as late as the thirty-fourth day, as did also that remaining in the bladder after death. Towards the end there was evidence of renal irritation. Experimentation has shown that after such a small dose of arsphenamine, arsenic does not usually appear for so long, or so consistently. The slight renal irritation that appeared just before the end cannot be ascribed to arsenic, but rather to the impairment of the skin function. It was only after large quantities of the scales had been tested that arsenic was discovered in them. There was a high polymorphonuclear leucocytosis, persisting up to within a day before death. There was a marked eosinophilia, as high as 40%, and the eosin cells were most numerous at a time when there was a slight improvement in the patient's condition, the percentage sinking to zero during the last three days of life.

The autopsy showed extensive passive congestion and an area of broncho-pneumonia at the right base. Heart and liver soft and pulpy, in many places fatty degenerated. Kidneys somewhat congested, but there was no definite nephritis. There was an adherent mucous on the mucosa of the stomach and intestines, which showed chemically the presence of arsenic, as did the liver, brain, skin, and the pericardial fluid.

In his summary, the writer emphasizes the fact that diarrhea and vomiting were absent during all stages of the intoxication, that nephritis was not marked at any time and appeared only at the end; that there was a decided affinity of the poison for the skin and for the trophic nerves supplying it; that arsenic was persistently present in the urine, without accompanying renal irritation, and was found at the autopsy in every tissue in which it was sought for; and that there was a high leucosytosis and eosinophilia, the latter corresponding closely to the patient's condition.

MORBID PHENOMENA FOLLOWING ARSPHENAMINE AND NEOARSPHENAMINE.

Beeson[4] reports a case of polyneuritis plus dermatitis exfoliativa following neoarsphenamine, an association the rarity of which is shown by a review of the literature. It was first mentioned by Duhot in 1912, who reported that in patients treated with large doses of neoarsphenamine, repeated every two or three days until four or five injections had been given, there appeared first, formication and pain in the soles and calves, edema of the legs and feet, and a reddening and furfuraceous scaling of the skin, followed in the more serious cases by successive disappearance of the plantar, Achilles, and patellar reflexes, together with loss of muscular power in the legs, and paralysis. There were similar symptoms in the arms. These cases were believed to be due to chronic arsenical intoxication. There were no cerebral complications, or disturbances of sight or hearing. Spillman has reported one similar case.

Beeson refers to the fatal cases of dermatitis exfoliativa after arsphenamine treatment recently reported by Latham and Carr and to a personal communication of an analogous case from Fordyce, Ormsby and Morrow, and remarks that it appears that exfoliative dermatitis has been observed by a large number of those who have had a wide experience with this treatment. Brooks and Roberts have described symptoms from an epidemic of arsenical beer poisoning in England in 1901, that correspond closely to those following arsphenamine and neo-arsphenamine poisoning, namely exfoliation of the skin, pigmentation, hyperkeratoses, pemphigoid eruption and herpes, and others have emphasized the nervous phenomena exhibited in this epidemic.

Beeson cites a number of other cases of polyneuritis following arsphenamine and neo-arsphenamine, and then proceeds to collate the various reported cases of peripheral nerve involvement due to syphilis and mercury. Of the former there are numerous instances, and a few well-recorded cases of the latter.

Beeson's own case is that of a negro of 42, who had a positive Wassermann, although there was no other evidence of syphilis. He had six weekly intravenous injections of neo-arsphenamine given in concentrated solution, and a dozen intramuscular injections of 1 per cent. solution of mercuric chloride. A second course was begun after three weeks, and two more injections of the neo-arsphenamine given, when he was taken suddenly ill and soon developed a tense and dangerous exfoliative dermatitis, attended with emaciation and delirium. Two months later, nerve symptoms appeared, and he soon developed a polyneuritis. Despite his serious condition and symptoms of marked toxemia, such as a high fever lasting almost three

months, he eventually recovered. The treatment was symptomatic.

It is considered that there is no reason to doubt that neo-arsphenamine was the causative factor, on account of its arsenical properties, and a cumulative effect is to be supposed. Beeson considers that syphilis and alcohol were contributing causes. He considers it improbable that mercury was a factor, since in that case we should expect stomatitis, more marked gastro-intestinal symptoms, and probably neuritis; also most cases of intoxication after mercurial injections come after the use of insoluble salts, such as calomel and gray oil.

ETIOLOGY OF ALOPECIA AREATA.[5]

Sabouraud, after more than twenty years' study of this mysterious affection, agrees with Brocq in the strength of the arguments in favor of local parasitism as the cause of the circinate form of alopecia areata and of its spreading out from a center. This view is supported by the fact that the area appears suddenly by the spontaneous falling of forty to fifty hairs, and increases progressively by the falling of hairs in the periphery. Among the hairs that fall are noticed the club-shaped or exclamation-point hairs, which have been described by Sabouraud as "cheveux peladiques," and which he regards as absolutely characteristic of plaques that are in process of extension; as when a primary plaque is checked in its course, one often sees groups of exclamation-point hairs at the periphery which are the forerunners of a new point of extension. Oftentimes, also, the areas may increase in one diameter while they fill in in another with new hairs, so that the area will end by healing at a very different point from that from which it started. Sometimes a single area may constitute the whole affection; it heals and everything disappears. At other times, however, perhaps six weeks after the appearance of the primary area, a series of similar patches may appear, similar to the original in all respects. Later on, after a period which might be called incubation, the plaques multiply still more. All this is similar to what happens in trichophytosis. Another point observed recently by Sabouraud is that after a primary plaque has been followed by a period of apparent repose, one often sees several secondary plaques appear, which are in turn followed, after several weeks more, by a multitude of new and very small plaques each of which, before becoming bald, is signalized by an erythematous spot of orange red, of the same dimensions as the succeeding area. This erythema is very transient, and is a rare occurrence.

In Sabouraud's opinion, the patch of alopecia areata, with its fringe of classic hairs, is a morbid type of special, peculiar characteristics. The only other condition that can be compared to it is that produced by the local action of the x-rays.

Another type of alopecia areata is that called ophiasis, which usually begins at the lower part of the occiput, extending upward, usually by way of the borders. Sabouraud considers that the two types differ only by symptoms which are not always present, while the essential symptoms are the same in the two types. In neither type is there the slightest proof of contagiousness, in Sabouraud's view.

The writer is not disposed to regard the evidence in favor of the trophoneurotic origin of alopecia areata as convincing, and combats one by one the various arguments that have been adduced.

As to the theory of the dental origin of alopecia areata, introduced by Jacquet, Sabouraud considers that a relationship between the teeth and this affection has not been shown. But in examining the teeth of his patients, he has been struck wit hthe frequency of multiple affections of the bones, such as deformities of the cranium, changes in the maxillary bones, etc. He points out that syphilis is considered to be the principal cause of these defects, although he does not claim that syphilis, either late or hereditary, is the sole cause of alopecia areata. Sabouraud considers it more important to treat the healthy portions of the scalp than the diseased. If the whole scalp is treated, the alopecia may, in many cases, be limited. Treatment of the whole head with oil of cade has given excellent results. Topically, he considers chrysophanic acid diluted 1-100 in chloroform the most active remedy.

CREEPING DISEASE.[6]

Darier has observed, and described with his usual painstaking accuracy, a case of creeping disease (larva migrans), an affection rarely seen in France, although zoölogists report that species of the organisms which cause it are common all over Europe and in Northern Africa. The clinical aspects of the form named

creeping eruption by Robert Lee in 1874, are quite characteristic. The case reported was at once recognized by all the assistants.

The patient was a man of thirty-nine, employed on the railroad in Paris. It was pretty conclusively shown that he must have contracted this parasitic affection in Paris. The lesion was situated on the back of the left hand, extending from the wrist to the lower part of the little finger, and took the form of a red and elevated streak, quite sinuous, and of an oedematous rather than indurated consistency. It had begun five days previously by intense pruritus. The parasite was extirpated by incising freely the tissues which contained it. It was found microscopically to be situated immediately under the epidermis of the finger, surrounded by a chitinous membrane armed with hooks, provided with a digestive tube, and muscular fibres in its body. There was a total absence of inflammatory reaction of the skin. The canal dug in the skin by the parasite by means, no doubt, of its hooks, was entirely filled by it and was nowhere below the skin, but immediately beneath the epidermis, in the papillary layer. It was thought that the larva might be that of the Hypoderma bovis of Geer, with which the patient may have come in contact through his employment at the Gare Montparnasse. Similar cases have been observed in the Shetland Islands, Norway, Brittany, Brazil, etc. Nowadays three forms of cutaneous and subcutaneous myiasis are recognized clinically: (1) Creeping disease, or myiasis rampante, characterized by a red sinuous line, raised and progressive, which advances from one to four centimeters per day, of which the above case is an example. (2) Subcutaneous myiasis, *à tumeurs ambulatoires,* in which oedematous or inflammatory tumors are produced during several weeks or months, which may attain the size of an egg, and the last one of which gives issue to the larva. The tumors are due to the subcutaneous migration of a larva which causes sharp pain. (3) Furuncular myiasis, in which there are no intradermic or subcutaneous migrations of the larva; a single inflammatory tumor is the first and last morbid manifestation, the parasite itself finishing by leaving the tumor, if it is not extracted. It is remarkable that each of these three clinical forms is not due to a different parasitic species, as larvae of diverse species have been found in each of these classes.

REFERENCES.

[1] Paris Médical, June 7, 1919.
[2] The Lancet, French Supplement, Sept. 27, 1919.
[3] Jour. A. M. A., July 5, 1919.
[4] Archives of Derm. and Syph., September, 1920.
[5] Annales de Derm. et de Syph., vi serie, tome 1, No. 4
[6] Annales de Derm. et de Syph., vi serie, tome 1, No. 3.

Book Review.

Tropical Ophthalmology. By Robert Henry Elliot, M.D., B.S., F.R.C.S., London: University Press, 1920.

No one who takes up this book will want to put it down until it is finished. It is attractive in style and undertakes a new subject. The scenes and conditions which have been familiar to Colonel Elliot awaken in us the liveliest curiosity, and his description of the natives in India and their environment, as far as it relates to this subject, is fascinating. A humane and conscientious surgeon, he dwells on the human side of his work and leaves a striking picture of medical work in the tropics. After discussing the mental and physical characteristics of the natives and the conditions which cause or favor ophthalmic disease, he depicts the peculiarities of these diseases as found in the tropics, giving special attention to injuries inflicted on the eyes by animals and parasitic diseases. These chapters are a mine of information. The chapters on operative work, especially cataract, and the work of the native coucher, will attract attention. The last section, dealing with general disease as affecting the eyes, gives the author a chance to express valuable views on some subjects we know little about in this climate, and his remarks on malaria as a cause of corneal condition are instructive. The book is well illustrated in black and white and has seven fine colored plates. We advise every ophthalmologist to read this book, not only for instruction, but also for pleasure.

Yale Medical School.—Dr. Yandell Henderson, hitherto professor of physiology in the Yale Medical School, has been transferred to the Graduate School of Yale University, under the title of Professor of Applied Physiology.

The tuition fee in the Medical School, beginning with the next university year, has been increased from $240 to $300, and announcement is made that, beginning with a year from next fall, three years of collegiate work or its equivalent, instead of the present two years, shall be required for all candidates for admission to the School of Medicine.

THE BOSTON
Medical and Surgical Journal

Established in 1812

An independently owned Journal of Medicine and Surgery, published weekly under the direction of the Editors and an Advisory Committee, by the Boston Medical and Surgical Journal Society, Inc.

THURSDAY, APRIL 7, 1921

Editors

Robert M. Green, M.D., *Editor-in-Chief and Manager*
George G. Smith, M.D., *Assistant Editor*

Advisory Board

Edward C. Streeter, M.D., Boston, *Chairman*
Walter P. Bowers, M.D., Clinton
Walter B. Cannon, M.D., Cambridge
Harvey Cushing, M.D., Brookline
Homer Gage, M.D., Worcester
Reid Hunt, M.D., Boston
Lyman A. Jones, M.D., Swampscott
Roger I. Lee, M.D, Cambridge
Robert B. Osgood, M.D., Boston
Milton J. Rosenau, M.D., Brookline
Alfred Worcester, M.D., Waltham

Subscription Terms: *$6.00 per year, in advance, postage paid for the United States, $7.56 per year for all foreign countries belonging to the Postal Union.*

An editor will be in the editorial office daily, except Sunday, from twelve to one p.m.

Papers for publication, and all other communications for the Editorial Department, should be addressed to the Editor, 126 Massachusetts Ave., Boston 17. Notices and other material for the editorial pages must be received not later than noon on the Saturday preceding the date of publication. Orders for reprints must be returned in writing to the printer with the galley proof of papers. The Journal will furnish free to the author, upon his written request, one hundred eight-page reprints, without covers, or the equivalent in pages in the case of articles of greater length.

The Journal does not hold itself responsible for any opinions or sentiments advanced by any contributor in any article published in its columns.

All letters containing business communications, or referring to the publication, subscription or advertising department of the Journal should be addressed to

Boston Medical and Surgical Journal
126 Massachusetts Ave., Cor. Boylston St., Boston 17, Massachusetts

ENCEPHALITIS WITH EXCITATION

Our attention has been called to the peculiar form of encephalitis, reported as "talking sickness," from which little eight-year-old Miriam Rubin, Waukegan, Illinois, suffers. This case has been the subject of a widely circulated newspaper story, based on half-truths, and commercialized by a widespread advertising propaganda. The newspapers have been deceived and misled. The public should know the facts ascertained by medical investigation. It is suggested that medical societies, particularly in localities in which the story has been circulated, should hold a meeting, with the object of appointing a committee to call on the local press. Where this is impossible without delay—for the action should be immediate since the story is a "live" one—the officers of the society might act, although a committee, of not less than three, would be more influential than an individual.

Undoubtedly the editors of newspapers will gladly coöperate with the local profession if the coöperation is requested in the right spirit. It would not be advisable, and nothing would be accomplished, to make the appeal in a spirit of anger or antagonism. The press should be approached realizing its viewpoint, which naturally differs from that of the medical profession, and should be asked that the following facts be given to the public:

1. That the child suffered, not from a strange "talking" sickness, but from a form of encephalitis with excitation.

2. The "incessant" talking was, in reality, intermittent, both before and after the alleged "adjustment."

3. The chiropractic "treatments" did not "cure" the disease, as claimed; they had no appreciable effect on its course.

4. The nurse's record shows that the chiropractor gave "treatments" from Februrary 12 to February 23, at which time he was dismissed, as the patient's condition gave no evidence of benefit from his "treatments." On the contrary, she was complaining of severe pain along the course of the spine. Since then, the family physician has had entire charge of the case.

5. The patient is not restored to health: on March 1, she still was seriously ill.

The medical profession is warranted in asking the press to give to the public the truth.

LICENSING OF PRIVATE HOSPITALS.

On March 16, there was held at the State House, by the Legislative Committee on Public Health, a hearing on Senate Bill number 2501, relative to the licensing of hospitals in Greater Boston. This proposed law provides that no license shall be granted for the maintenance of a hospital, if the property owner within one hundred feet of the hospital objects. Under the definition of a hospital, this law might include every doctor's office where sick patients are cared for, and every convalescent home or sanatorium. Obviously, this situation would be productive of injustice and hardship. The private hospitals of Boston have organized to meet the emergency and will endeavor to secure an unfavorable report upon the bill. It is hoped that physicians who do not desire its passage will assist the hospitals with such influence as they may possess.

MEDICAL NOTES.

WISCONSIN HOME-COMING.—The State Medical Society of Wisconsin will celebrate its seventy-fifth birthday by holding a "Home-Coming" meeting in Milwaukee, September 7, 8 and 9, 1921. All former Wisconsin men, whether they have practiced there or left Wisconsin to study medicine, practising elsewhere after graduating, are invited to this home-coming.

The officers of the Society are anxious to secure at this time, for mailing purposes, the names of all former Wisconsin men. They will confer a favor by sending their names and addresses to Dr. Rock Sleyster, Secretary, Wauwatoca, Wisconsin.

APPOINTMENT OF DR. C. L. ALSBERG.—Dr. C. L. Alsberg, Chief of the Bureau of Chemistry of the United States Department of Agriculture, has been appointed Director of the Food Research Institute, which is to be established at Stanford University by the Carnegie Corporation. He will assume his new work on July 1.

BOSTON AND MASSACHUSETTS.

WEEK'S DEATH RATE IN BOSTON.—During the week ending March 26, 1921, the number of deaths reported was 211 against 268 last year, with a rate of 14.53 against 17.28 last year. There were 39 deaths under one year of age against 40 last year.

The number of cases of principal reportable diseases were: Diphtheria, 67; scarlet fever, 72; measles, 93; whooping cough, 27; typhoid fever, 2; tuberculosis, 49.

Included in the above, were the following cases of non-residents: Diphtheria, 4; scarlet fever, 5; typhoid fever, 1; tuberculosis, 10.

Total deaths from these diseases were: Scarlet fever, 5; whooping cough, 2; tuberculosis, 15.

Included in the above, were the following cases of non-residents: Tuberculosis, 2.

EXHIBIT OF RARE OLD BOOKS ON FEVERS AND SPECIFIC INFECTIONS AT THE BOSTON PUBLIC LIBRARY.—During the meeting of the American Medical Association here in June, in the room used for exhibits on the floor of the Special Libraries at the Public Library, there will be spread an exhibit of early texts (Hippocrates to Sydenham) dealing with fevers and with the specific infections, arranged in chronological order. Besides the texts, there will be considerable illustrative material touching hospitalization and treatment, the use of baths, venesection, new remedies, pest banners, broadsides and medals, also Saint Roch and Saint Sebastian, and various aspects of the Plague and Syphilis dealt with in the graphic arts. General tests illustrating the Greek, Byzantine, Mohammedan and medieval practice in fevers will occupy one half of the space allotted, while the other half will contain tracts on the Plague and Syphilis, original descriptions, new diseases and primary treatises on the doctrine of *Contagium vivum*. A descriptive catalogue will be ready in time for the exhibit, which will last from June 5 to 11.

APPOINTMENT OF DR. WILLIAM A. BRYAN.—Dr. William A. Bryan has been appointed superintendent of the Worcester State Hospital. Dr. Bryan has been assistant superintendent of the Danvers State Hospital, before which he served on the staff of the Boston Psychopathic Hospital. Dr. Bryan will succeed Dr. Ernest V. Scribner.

RETIREMENT OF DR. EDWARD L. MARK.—Dr. Edward Laurens Mark, who has been instructor and professor of zoölogy and anatomy at Harvard University for forty-four years, will retire from active teaching at the close of the present year. He has been appointed Hersey professor of anatomy emeritus.

NEW ENGLAND NOTES.

OBSERVANCE OF DR. STORER'S BIRTHDAY.—Dr. Horatio R. Storer, dean of the medical profession of Newport, R. I., and in former years active in practice and in various scientific and health protective organizations, was honored on the occasion of his ninety-first birthday, March 20, when he was presented by representatives of citizens of Newport with a silver loving cup, in recognition of his services to the community and as an expression of the esteem in which he is held. During the day, he received many callers at his home, and many gifts and messages from friends in this city and other

parts of the world. Among those whom he had with him on that day were his two sons, Dr. Malcolm Storer and Mr. John Humphrey Storer, of Boston.

During the early part of the day, Dr. Norman M. McLeod, Dr. Edward V. Murphy and Chaplain E. S. Burke of the Training Station, called at the Storer residence and, as representatives of the donors, presented the loving cup. Dr. McLeod made the presentation in a brief speech, expressing the good wishes of the donors. Dr. Storer showed his appreciation of the testimonial in a brief reply, in which he said he felt that the gift was not so much a tribute to himself as the work which he, with the donors, had been able to do for the best interests of others.

The cup, which is of beautiful design, is inscribed as follows:

Horatio Robinson Storer
the Physician
Pioneer Surgeon, Philanthropist,
Leader in Civic Administration
and the
Always Helpful Adviser,
From His Admiring Fellow
Citizens and Friends of
Newport, Rhode Island.
February 27th, 1921.

The donors included the physicians of the city, the Redwood Library, the Newport Historical Society, the Advisory Council of the Civic League, the Harvard Graduates of the city, several Roman Catholic Church organizations, and many individual friends.

Correspondence.

ACTION TAKEN BY THE PROFESSIONAL GUILD OF KING'S COUNTY.

New York City, March 17, 1921.

Mr. Editor:—

Physicians and druggists everywhere, throughout the United States, will be interested in the action of the Professional Guild of King's County in deciding to ask for an injunction restraining the New York State Narcotic Drug Commission from enforcing recent rulings of the department, compelling the use of official triplicate blanks for all prescriptions containing narcotic drugs. This course was decided upon at a meetng just held in Brooklyn, which was largely attended by the doctors, druggists and dentists, who make up the membership of the Professional Guild. At present, this ruling makes the doctors of the state, technical violators of the law, at least, as few of them have provided themselves with these blanks, believing that, under a proper construction of the Harrison Law, they are permitted to use the non-official blanks in their daily practice.

It is the general consensus of opinion that the Professional Guild is entitled to great credit for its action in this matter. Properly speaking, this course should have been taken by the organized medical bodies of the state, which, however, with one exception, have permitted this ruling to go uncontested. This is true, also, in a measure, of the state officials responsible for the acts of a subordinate official who, under the law, is empowered to make needful and helpful regulations in the conduct of this office. Needless to say, such regulations would not be universally disregarded by a class of citizens represented by the physicians and druggists of this state, if they did not endanger the safety and lives of those whom they are bound to protect, even at the risk of their own safety and welfare. This inaction cannot be excused, either, by the fact that bills are now before the Legislature abolishing the office of Narcotic Drug Commissioner.

Even if this be the case, such conduct should not be condoned indefinitely until such time as the offender is legislated out of office for reasons of economy only, when there ars many more substantial reasons for taking action upon the administration of this office at the present time.

John P. Davin, *Executive Secretary,*
New York Medical Association.

THE DANGER OF TYPHUS.

Mr. Editor:—

Is this country confronted with an epidemic of typhus, brought about by the flow of immigration, or the danger nil?

The public has full right to await an answer from, and expect all precautions to be taken by, medical men, whose duty it is to protect the population from an epidemic that is preventable. Having recently returned from Europe, where typhus is raging, I believe that it will be of interest if I relate a few of my experiences on this subject.

The people of Russia and Poland, with a few exceptions, are unfortunate in not having any baths in their homes, and they are obliged, therefore, to use public baths. Recently, the absence of fuel has caused the closing down of these baths. This condition, coupled with the scarcity of soap, water and underwear, resulted in conditions where vermin are multiplying without interference.

Every medical man knows how rapidly lice multiply, and thus, in the deplorable condition, I found Russia infested with vermin. A few hundred of these oviparous insects found on the human body is considered an ordinary case. I remember, when making calls on patients in Russia, I would place a chair in the middle of the room and leave my coat and hat on it, telling the people to keep away from the chair. Arriving at home, I would inspect my clothing and would invariably find three or four of the vermin.

The lice have so changed their adaptations to surroundings that they live on the walls, furniture, etc. The wooden walls and ceilings of the freight cars, which are now used for passengers, are infested with these vermin.

I can also remember that later, while in Constantinople, on the Island of Proti, where the American Near East Relief Committee took care of the Russian refugees, I warned Major Arnold, the director, to be careful while in contact with the Russians on the ship that brought them to the island. He smiled, thinking it to be imaginary. In his great desire to help these unfortunate people, he forgot the danger, and I constantly had to remind him by drawing his attention to the thousands of lice "promenading" on the clothes of the refugees.

Living in unsanitary conditions and being very poorly nourished, is it a wonder that in the city of

Odessa alone, there are over three hundred deaths daily, caused by typhus?

A station near Proskurff was on the battlefield between the Soviet and Ukrainian troops, but none dared to pass it, for it was full of typhus victims, both dead and alive. The two warring armies stood on both sides of the station for a long time, fearing to cross the typhus line.

The Germans have tried to impress upon the population of Constantinople, during the war, the cause of typhus, by putting out coffins at certain corners of streets and having a photograph of a large louse stinging the forehead of a man, who falls into the coffin. Thus typhus was named by the people "louse fever."

The prevention of the spread of the disease should be done thoroughly, no break being allowed anywhere, and, unless carried out properly, the disease will increase and multiply.

Very energetic measures were taken by us to stamp out typhus among the refugees on the Island of Proti (the American Island), letting the whole colony of Russians through a bath every three or four days, delousing their bedding, belongings, etc., at the same time, and only the rigid discipline on our island, and the strict businesslike manner in which we conducted the work, resulted in checking the disease with a small mortality, while the British, who were rather lax, had a high mortality, losing also one of their own physicians from typhus.

I am proud to state that, in spite of the tremendous work done by the American staff, and the constant contact with typhus patients, no one contracted the disease, while in one of our hospitals (I B) the Russian physician and all the Russian nurses contracted typhus, due to loose conduct.

Before leaving the island, every refugee was obliged to take a bath and allow all his things to be put through the delouser. An inspection of the refugees was then made and a certificate of disinfection was given when he was found free of vermin. But, after taking the bath, over fifty per cent. were found lousy, and frequently the same procedure was repeated three or four times before he was found free from vermin.

We learned that taking a bath does not always clean up a lousy person.

On my way to the States, I naturally was quite interested in finding out what the American authorities do to prevent bringing the typhus epidemic into the States, so in Patras (a large Greek port for immigrants) I went to the American Consul to consult with him regarding the precautions taken before the immigrants were allowed to leave the city. He assured me that everything was done according to science; every passenger was obliged to take a bath and his baggage was disinfected. My heart was full of joy to see the big signs on the immigrants' trunks, "disinfected," and my joy was coupled with merriment when I saw, outside the ticket office, the passengers busying themselves pasting on their trunks, which they had just brought from the hotels, the labels "disinfected," just received for a fee from the office.

I do not know what science was required to paste these labels on, but the consul was quite satisfied that it meant medical science.

In conclusion, I wish to state that I do not want to be quoted as saying that the fear of typhus should result in stopping immigration—it would be an insult to the medical profession—but I do believe that we should look upon the question of bringing into this country an epidemic of typhus very seriously, and that we should handle the matter with due consideration, making no distinction of the passenger, be he immigrant or American, first class or steerage.

S. G. PAVLO, M.D.
356 Ferry Street, Malden, Mass.

SHARON SANATORIUM NOT A STATE INSTITUTION.

Boston, March 31, 1921.

Mr. Editor:—

The directors of the Sharon Sanatorium are deeply grateful to you for the admirable editorial in the edition of Thursday, March thirty-first. There is a mistake, however, in the early part of the article which we feel it is necessary to correct, as it may give a wrong impression to your readers.

You say: "The step taken by Massachusetts in establishing the Sharon Sanatorium as a *state* institution is particularly significant," etc. The Sharon Sanatorium has never had anything to do with the state. It was founded by private subscription, and has been largely supported by private subscription ever since, and has never received any money from the state. It was the first institution of its kind in New England, and it was an experiment to prove that much more could be done than hitherto had been thought possible in the cure of tuberculosis at a low altitude and in harsh climates like that of New England. The first state sanatorium in the Union was opened at Rutland, Massachusetts, in 1898, seven years after the establishment of the Sharon Sanatorium. In the preliminary report of the later institution, the trustees stated that it was the success of the work at Sharon that induced them to completely change the character of the institution at Rutland, this change being due chiefly to the efforts of Dr. Alfred Worcester of Waltham, Massachusetts, one of the Trustees, who had watched the work at Sharon and believed that the character of the state institution should be changed from that of a mere hospital for consumptives, to a sanatorium for hopeful cases.

As the directors are particularly desirous that the means of support of the Sharon Sanatorium should be known to the public, and that it is in no way connected with the state, I trust you will be willing to print this correction in your next edition, and by so doing greatly oblige us.

With renewed thanks for the editorial, I remain, in behalf of the directors,

Yours very truly,

VINCENT Y. BOWDITCH.

SOCIETY NOTICES.

MASSACHUSETTS SOCIETY FOR MENTAL HYGIENE.—The Human Element in Industry is the subject of the Annual Conference of the Massachusetts Society for Mental Hygiene to be held on Thursday, April 7, at 8 P.M., at Ford Hall, Boston. Dean Everett W. Lord of the College of Business Administration, Boston University, is to preside, and the following is the program:

1. "Has Mental Hygiene a Place in Practical Industry?" Boyd Fisher, Service Department, Lockwood, Greene & Co., Boston.
2. "Industrial Hygiene." Wade Wright, M.D., Secretary of the Division of Industrial Hygiene, Harvard Medical School.
3. "The Mental Health of the Industrial Worker." C. Macfie Campbell, M.D., Director of the Boston Psychopathic Hospital.

The public is cordially invited to attend this meeting.

SUFFOLK DISTRICT MEDICAL SOCIETY.—Censors' Meeting. The Censors of the Suffolk District Medical Society will meet for the examination of candidates at the Medical Library, No. 8 The Fenway, Thursday, May 5, 1921, at four o'clock.

Candidates should make personal application to the Secretary and present their medical diploma at least one week before the examination.

R. H. MILLER, M.D., *Secretary*,
402 Marlborough St., Boston.

PURITY
SAFETY
ENDOARSAN
(Syphilis)
Dimethylarsenate Mercury
Iodides, Phosphorus
ENDOMERSAN
(Cerebro-Spinal Syphilis)
Mercury, Dimethylarsenate
Iodides
"ENDO"
Specialties
for Direct Medication
by the
Intravenous Method
ENDO-QUIN
(Malarias)
Quinine Hydrochloride
ENDO-GLOBIN
(Chlorosis Pernicious Anemia)
Haemoglobin
To the modern physician, awake to the advantages of Direct Medication, we offer a complete line of pure and stable products which may be injected into the blood stream with the certainty of no untoward effects.
Our Intravenous Ampoules form the border of this announcement. Acquaint yourself with their names, ingredients and the conditions in which they are indicated.
Do not confuse us with any other firm of similar name, or with those making similar preparations. Insist on "ENDO" products, the formulae and mode of adminstration of which insure the best results in:
Syphilis Malaria Toxemias
Tuberculosis Asthma Cystitis
Rheumatism Anemias Pyelitis
Bronchial and Streptococcus Infections
Ask for catalogue giving complete formulae of our specialties. Reprints of interesting articles and price list will accompany it. Correspodence is invited and will be promptly replied to by one of the physicians on our staff.
ENDO-CREODIN
(Bronchial and Pulmonary Affections)
Iodides, Guaiacol, Creosote
ENDO-SAL
(Rheumatism, Streptococcus Infections)
Iodides Salicylates Colchicine
Intravenous Products Co. of America, Inc.,
121 Madison Ave., New York City
Most Effective Method in the Treatment of Disease
ENDO-SODIUM IODIDE
(Asthma)
2 Grams of Sodium Iodide
ENDOFERARSAN
(Anemias)
Iron and Arsenic
ENDOMETHYLENAMIN
(Toxemias, Pyelitis, Cystitis)
0.6 Gram Hexamethylenamine
ENDO-CAODIN
(Tuberculosis)
Calcium Iodides Guaiacol
EFFICIENCY
RELIABILITY

900 BEACON ST.

To rent, entire second floor for doctor's office, with privilege of living in same.
10-tf,B390

FOR SALE

$5,000 country practice in extra good locality. Price $1,500, reasonable terms. Address
A276, Care of
Boston Medical and Surgical Journal
6-tf

A SMALL HOME SCHOOL

in the pine woods, for delicate or young children. Care of Trained Nurse. Excellent teaching. Outdoor life. Apply to
Rev. Mother, O. S. A.
St. Anne's House
44 Temple St., Boston, Mass.
2-tf,B363

PHYSICIAN'S OFFICE TO LET

All office conveniences, reception hall, good service. Near Kenmore Station. Inquire at
74 BAY STATE ROAD
6-tf,B377

TO LET

Offices for doctors or dentists. Single or in suite With service.
1069 BOYLSTON ST., ROOM 3
Tel. B. B. 59426 6-tf,B379

TO LET

At 587 Beacon St., Boston, junction of Bay State Road and Beacon St., two minutes from Kenmore Station, office overlooking Charles River. Large waiting room. Apply at 587 Beacon St., from 9 A.M. to 5 P.M. 7-tf,B381

PRACTICE FOR SALE

Unopposed practice in New Hampshire town of about 1000. Reasonable terms. Leaving to specialize. Address
A284, Care of
Boston Medical and Surgical Journal
13-tf

WANTED

Physician, male preferred, for Asst. Supt. at state sanatorium for tuberculosis, at Wallum Lake, R. I. Institution well equipped for the care and treatment of all stages of tuberculosis. Fine x-ray plant, staff discussion of all cases. Initial salary, $2000, with maintenance of self and family. Increases for one who makes good. Applicant should have had some tuberculosis experience. State age, height, weight, health, preliminary education, medical school, date of graduation, number in family, experience and references in first letter.
Dr. H. L. BARNES
Wallum Lake, R. I. 13-1,B393

TO LET

Desirable offices for physicians in newly remodeled building on Commonwealth Ave., at Mass. Twenty-four hour service. Apply to
A283, Care of
Boston Medical and Surgical Journal
12-tf

WANTED

Male physician at the Wrentham State School. Splendid opportunity to become familiar with the care and management of the feeble-minded. Applicants must be graduates of a Class A medical school, single, and not over 35 years old.
GEORGE L. WALLACE, Supt.
Wrentham, Mass.
15-tf,B396

WANTED

Electric Cabinet Bath; Massage Table and Victor Vibrator. Also Electric Wall Plate. Address
ALFRED WITTIG
15 Exchange St., Boston
Tel. Main 1798 14-2,B394

WANTED

Physician would like small office, furnished or unfurnished, with running water and electricity, in Back Bay district. Address
P. O. BOX 422, QUINCY, MASS.
14-1,B395

WANTED

Office, of a busy practitioner in or near Boston, who wishes to go away for summer, by energetic young physician (29 years) from June 15th to Sept. 1st to fill in between two hospital appointments. Address
A285, Care of
Boston Medical and Surgical Journal
14-1

WANTED

Male assistant physician, who has had insane hospital experience, to fill vacancy on the staff of a State Hospital for the Insane. Salary, $2,000 per year with maintenance. Address
A286, Care of
Boston Medical and Surgical Journal
14-tf

BIND YOUR VOLUMES

of the

BOSTON MEDICAL AND SURGICAL JOURNAL

We will furnish the covers, have them bound, and delivered to your office.

Half-year volumes at $1.75 each
Complete year in one volume, $2.00

THE BOSTON
Medical and Surgical
JOURNAL ..

126 Massachusetts Avenue

YOUNG'S HOSPITAL

788 MASS. AVE., ARLINGTON
Established 1910
Surgical Medical Obstetrical
OPEN TO THE PUBLIC
Physicians and Surgeons may secure accommodations for a limited number of free patients
SEND FOR CIRCULAR
R. D. YOUNG, M.D., Director
GEORGIA E. YOUNG, R.N., Supt.
FANNIE L. ROBBINS, R.N., Supervisor of Nurses
4-eow,1y

A Relief for Chronic Constipation

During Pregnancy

For Nursing Mothers

FRUTOSEN

The Laxative
Par Excellence
For Children

Palatable No Griping

In 4-ounce bottles

FRUTOSEN DRUG CO.
Boston, Mass.

W.S.S.
WAR SAVINGS STAMPS
ISSUED BY THE UNITED STATES GOVERNMENT

W.S.S.
WAR SAVINGS STAMPS
ISSUED BY THE UNITED STATES GOVERNMENT

WAPPLER

ELECTRO-MEDICAL APPARATUS

X-RAY MACHINES AND ACCESSORIES
HIGH FREQUENCY MACHINES
GALVANIC, FARADIC, SINUSOIDAL APPARATUS
REPAIRS—SERVICE—SUPPLIES

CHAS. H. KEHLENBACH, 1074 Boylston Street
SALES AGENT BOSTON, MASS.
TEL. BACK BAY 8116

BOURNEWOOD HOSPITA

SOUTH STREET, BROOKLINE, MASS.

Established 1884

For a limited number of cases of Mental and Nervous Disease only
Post Office, Chestnut Hill. Telephone, "Parkway" 300
Nearest station, Bellevue, N.Y., N.H., & H. R.R.

GEORGE H. TORNEY, M.D.
Physician in Charge

HENRY R. STEDMAN, M.D
Consultant

CHANNING SANITARIUM, Inc.

Wellesley Avenue, Wellesley, Mass.

(Established in Brookline, 1879. Transferred to Wellesley, 1916.)

Seven new buildings on fifty acres of high woodland. Sleeping porch and private bath for each patient. Large and small suite cottages. Separate buildings for men and women. Facilities for occupation and diversion. Complete equipment for Vichy, Nauheim and electric baths and other forms of hydrotherapy. Alpine lamp for heliotherapy.

Especially designed and equipped for the care and treatment of nervous cases.

DONALD GREGG, M.D. WALTER CHANNING, M.D.

GONOSAN
"RIEDEL"

To ALLEVIATE PAIN, to PROMOTE DIURESIS and to PROTECT the membrane of the urethra, especially THE POSTERIOR PORTION—these are the important objects of the treatment of acute cases of Gonorrhea.

The entire urinary tract should be influenced by means of proper internal medication. Local injections alone will not be sufficient.

This is the rationale of GONOSAN.

Samples and literature at your disposal.

RIEDEL & CO., Inc.

Bush Terminal Building 5 BROOKLYN, NEW YORK CITY

THE STORM BINDER AND ABDOMINAL SUPPORTER

(PATENTED)

Adapted to use of Men, Women, Children and Babies

Modifications for Hernia, Relaxed Sacro-Iliac Articulations, Floating Kidney, High and Low Operations, Ptosis, Obesity, Pregnancy, Pertussis, etc.

Send for new folder and testimonials of physicians. Mail orders filled at Philadelphia only—within twenty-four hours.

KATHERINE L. STORM, M.D., 1701 Diamond Street, **Philadelphia**

Professional Cards

RING SANATORIUM
AND
HOSPITAL
Inc.

Arlington Heights, Massachusetts

Eight Miles from Boston
Telephone, Arlington 81

For Chronic, Nervous and Mild Mental Illnesses. Five separate houses permit of segregation and a variety of accommodations.

Boston Office—By appointment
496 Commonwealth Ave.
Telephone, Back Bay 8774
ARTHUR H. RING, M.D.,
Superintendent.

Dr. Mellus' Private Hospital
FOR MENTAL DISEASES
419 Waverley Avenue, Newton, Mass.
Reached by train to Newton, or by electric cars via Commonwealth Avenue to Grant Avenue.
Edward Mellus, M.D.

DR. TAYLOR'S PRIVATE HOSPITAL
For the Treatment of
NERVOUS DISEASES
ALCOHOLISM AND DRUG ADDICTIONS
House newly equipped and furnished, skilled attendants, good food, and comfortable rooms at moderate rates.
Methods of treatment are those proved best after 17 years' successful experience.
FREDERICK L. TAYLOR, M.D.
45 Centre Street, Boston
(Near Eliot Square) (Roxbury District)

MRS. SPINNEY'S HOUSE
A restful home for invalids and elderly persons, especially for the building up of convalescents amid attractive surroundings. Spacious piazzas, well stocked library, trained attendants.
MARY E. H. SPINNEY
17 Parley Vale, Jamaica Plain, Mass.
Tel. Jamaica 2230

"BELLEVUE"
Superior homelike accommodations for five patients. Nervous and Mild Mental Disease, selected cases of Alcoholism, and Elderly Women for whom medical supervision is desired, are received.
MARY W. L. JOHNSON, M.D.
158 WALCOTT ROAD, CHESTNUT HILL, MASS.
Telephone, Brookline 5381-W

WOODLAWN SANITARIUM
FOR THE CARE OF EPILEPTICS
DR. H. W. HAMMOND, *Supt.*
WEST NEWTON, MASS.
tf

Wiswall Sanatorium
Inc.
A Sanatorium for the Treatment of Nervous and Mental Diseases
New Buildings, every facility for comfort; in midst of twelve acres of high land, covered with beautiful oak and pine trees, fully equipped for hydrotherapeutic and electrical treatment.
H. O. SPALDING, M.D.,
Superintendent
E. H. WISWALL, M.D.,
Asst. Physician
Wellesley, Mass.
Telephone, Wellesley 261

DR. GIVENS' SANITARIUM
(THE STAMFORD HALL COMPANY)
STAMFORD, CONN.
For the scientific treatment of Nervous and Mental diseases, Drug and Alcoholic addictions, and General Invalidism. Completely equipped for the care and comfort of patients—Hydro-Therapy, Electro-Therapy, Massage, Occupational Therapy and Amusements.
Located in a beautiful natural park of 100 acres, with numerous detached buildings, insuring privacy.
Fifty minutes from New York City on the New Haven Railroad.
Address
Frank W. Robertson, M.D., Pres. and Med. Dir.
Phone. 70 Stamford
New York City Office: 412 West End Ave.
Monday, Wednesday, and Friday at noon.

WOODSIDE COTTAGES
FRAMINGHAM, MASSACHUSETTS
BOSTON OFFICE, 178 COMMONWEALTH AVE.
TEL. BACK BAY 808
A small sanitarium, pleasantly located on high ground, especially adapted to the care of chronic invalidism, rest cases, convalescents, and the aged. Separate groups. Committed cases not received.
Tel. Framingham 26.
FRANK W. PATCH, M.D.

GLENSIDE
For Nervous and Mental Diseases
6 Parley Vale.
Jamaica Plain, Mass.
MABEL D. ORDWAY, M.D.
Telephone, Jamaica 44.

HERBERT HALL HOSPITAL, Inc.
WORCESTER, MASS. Established in 1872

A Hospital for the Care and Treatment of those afflicted with the various forms of Nervous and Mental Diseases. Accommodations for a few chronic quiet patients at special rates.
WALTER C. HAVILAND, M.D., *Superintendent*
H. LINCOLN CHASE, M.D., *Resident Physician*

Attleboro Springs
A place for concentrated rest, convalescence from operations or prolonged illness, treatment of nervous and chronic diseases and nutritional disturbances. Mental and tubercular cases not received.
Regulated exercise, massage, hydrotherapy, electrical treatments. Diet scientifically supervised.
Modern building, airy rooms, 100 acres beautiful park and woodlands, spring water, farm and dairy.
Resident doctors and nurses.
Illustrated booklet sent on request.
ATTLEBORO SPRINGS
ATTLEBORO, MASS.
Tel. Attleboro 869.

THE MILLET SANATORIUM
FOR TUBERCULOSIS
EAST BRIDGEWATER, MASSACHUSETTS
C. S. MILLET, M.D., MEDICAL DIRECTOR
The Boston office of the Millet Sanatorium, East Bridgewater, Massachusetts, has been removed to
45 BAY STATE ROAD
Dr. Millet may be seen at this office on Tuesdays and Fridays between the hours of 11 and 1 o'clock.

TUFTS COLLEGE MEDICAL AND DENTAL SCHOOLS

The Tufts College Medical School offers a four-year course leading to the degree of Doctor of Medicine. The next session begins September 19, 1921. Students of both sexes are admitted upon presentation of an approved high school certificate and, in addition, college credit indicating two years' work in Chemistry, English, Physics, Biology, and French or German.

CHARLES F. PAINTER, M.D., *Dean.*

Beginning January, 1921, all candidates for admission to the Tufts College Dental School must present evidence of having attended one year in a recognized college or university, and to have passed the subjects of the year. These subjects must include Physics, Chemistry, Biology and English.

Tufts College has arranged a One-Year Pre-Dental Course which fulfills these requirements. A special bulletin describing the course will be sent upon application.

WILLIAM RICE, D.M.D., *Dean.*

For further information, apply to FRANK E. HASKINS, M.D., *Secretary*, 416 Huntington Avenue, Boston 17, Mass.

THE NEW YORK EYE AND EAR INFIRMARY
School of Ophthalmology and Otology
For Graduates of Medicine

Clinics daily by the Surgical Staff of the Infirmary. Special courses in Ophthalmology, Refraction, Operative Surgery of the Eye and Ear, Pathology and External Diseases of the Eye.

The abundant clinical material of this well-known institution affords students an unusual opportunity for obtaining a practical knowledge of these special subjects. Two vacancies in the House Staff exist in March, July, and November of each year. For particulars address the Secretary,

DR. GEORGE S. DIXON, NEW YORK EYE AND EAR INFIRMARY.

HARVARD DENTAL SCHOOL
BOSTON, MASS.

A department of Harvard University

Send for Announcement

A four-years' course

DR. EUGENE H. SMITH, *Dean*

SYRACUSE UNIVERSITY
COLLEGE OF MEDICINE

ENTRANCE REQUIREMENTS: Two years in a registered College or School of Science. Combination courses recognized.

LABORATORY COURSES in well-equipped laboratories under full time teachers.

CLINICAL COURSES in the University Hospital, one general, one special, and in the municipal hospitals and in the dispensary adjoining the college, in all of which senior students serve as clinical clerks. Tuition $200.

Address The Secretary of the College of Medicine, 307 Orange St., Syracuse N. Y.

GRAND VIEW SANITARIUM AND HOSPITAL
547 Central Street, Winchendon, Mass.

A Modern Health Resort for Convalescent Cases, Semi-Invalids, and General Medical and Surgical Work. Beautifully located with views of Monadnock in the north and Wachusett in the south. Pine grove. Rates moderate.

Licensed Lying-in Hospital. Confinements a specialty.

Doctors' private patients will receive our special care.

Write for particulars.

Julia A. Nason, R.N., Supt. Howard N. Nason, M.D., Physician

BOSTON UNIVERSITY SCHOOL OF MEDICINE
Founded 1873

REQUIRES FOR ADMISSION a minimum of two pre-medical college years including Chemistry, Physics, Biology, and, in addition to English, at least one foreign language.

REQUIRES FOR GRADUATION a minimum of four years of graded medical studies.

THE SCHOOL OFFERS: 1. A graded five-years' course, including hospital internship; 2 A six-year combination course with the College of Liberal Arts of Boston University, whereby a successful candidate may obtain the degree Sc.B., and the M.D.; 3. Eligibility to the Graduate School of Boston University to courses leading to the degree of Ph.D.; 4. A diversified curriculum covering the entire range of modern medical practice, including physiologic and homeopathic materia medica and therapeutics; 5. Unusually extensive facilities in the Boston City Hospital, Massachusetts Homeopathic Hospital, Haynes Memorial for Contagious Diseases, Robinson Maternity, Evans Memorial for Clinical Research and Preventive Medicine, and the Westboro State Hospital for the Insane.

Information gladly furnished on application to the Registrar,

EDWARD E. ALLEN, M.D., 80 East Concord St., Boston.

BOSTON NURSES' CLUB, Inc., AND REGISTRY
539 BOYLSTON ST., BOSTON, MASS.

BACK BAY 8787
SWITCHBOARD WITH TWO TRUNK LINES

Graduate Nurse in attendance day and night.

Graduates, Undergraduates, Attendants, Hourly Nursing, Masseurs, Male Nurses.

MISS FLORA E. ROSS, R.N., REGISTRAR

WET NURSES NEEDED
to fill increased demand at
WET NURSE DIRECTORY

Under the direction and control of the Infants' Hospital. Wet nurses may be obtained by telephoning to

THE DIRECTORY, Brookline 2930

A POINT SCALE FOR THE MEASUREMENT OF INTELLIGENCE IN ADOLESCENT AND ADULT INDIVIDUALS

Robert M. Yerkes, Ph.D., Boston, and Cecilio S. Rossy, M.A., New York

A description of the examples and application of the Point Scale used in the Army and for other scientific work along these lines. Reprints may be secured from this office at

35 Cents per Copy

The Boston Medical and Surgical Journal
126 Massachusetts Avenue,
Boston 17, Mass.

"THE REST"
53 Mass. Ave., E. Lexington, Mass.

Convalescents, semi-invalids, and persons recovering from operations given excellent care under the supervision of a Graduate Nurse.

Mrs. MARY G. MULLEN, Matron
Tel. Lexington 51663

CLINICAL CHART OF RENAL DISEASES

By H. S. Jellalian, M.D., of Boston

Student and practitioner alike will appreciate this concise tabulation. For wall or desk. 15 x 10 inches. 25 cents.

Boston Medical and Surgical Journal
126 Massachusetts Ave., Boston, 17 Mass.

tf

BOYLSTON LABORATORY
for
PHYSICIANS AND SURGEONS
Suite 308, 419 Boylston St., Boston

G. B. Whall, Manager

All forms of analytical work promptly executed

Fresh cultures of the Bacillus Bulgaricus made daily

Tel. B. B. 4200

For DIABETES

The best starch-free foods
are made from

LISTERS DIABETIC FLOUR
(PREPARED CASEIN)

Warranted strictly
starch-free and sugar-free.

Especially useful in a strictly non-carbohydrate diet.

Listers Flour is self-rising and easily made into palatable bread and pastry that satisfy diabetic hunger.
A months supply (30 boxes) $4.85

Ask us what store in your city has it - or it may be sent direct.

Made by LISTER BROS. Inc. 405 Lexington Ave. New York N.Y.

"If you have a castor oil that is truly tasteless and 'sets well' on the stomach, I want to know it and will be glad to use it."

This sentiment, expressed by one physician, voices the attitude of the entire medical profession.

Kellogg's Tasteless Castor Oil is made tasteless by refinement, not by the addition of any flavoring or disguising substance.

Kellogg's Tasteless Castor Oil is really tasteless; in fact, it has been substituted for olive oil in mayonnaise dressing without detection!

Kellogg's Tasteless Castor Oil is specially prepared for medicinal use. It is not only possible but easy to administer this most valuable agent in the treatment of obstipation, stasis, intestinal toxemia, hemorrhoids, asthma, acute bronchitis, and other conditions for which it was recommended in the Papyrus Ebers and other ancient medical works, and by physicians from Galen and Dioscorides down.

A clinical test of *Kellogg's Tasteless Castor Oil* will demonstrate its supremacy over ordinary castor oil or any flavored or disguised product offered as a substitute.

Send for interesting booklet, "*Therapeutic Uses of Oleum Ricini*" and sample, if desired.

Sole Distributor

WALTER JANVIER, Inc., 417-421 Canal Street, New York, N. Y.

If It Is Electrotherapeutic Apparatus We Have It
and
If It Is The Best We Have It

We carry a full line of Electric Light Bath Cabinets and Bakers, High Frequency Machines, Sinusoidal, Galvanic, and Faradic Outfits, etc.

We can give you the LATEST and BEST in X-ray equipments of any size complete with COOLIDGE accessories.

To physicians in New England who are using the Van Houten and Ten Broeck Static Machines we can give full service both for repair work and new equipment, as we have taken over the manufacture of these machines.

If you are in the market for any equipment it will pay you to write us.

Thomas B. Snow

212 East 23rd Street New York City

Anusol
HEMORRHOIDAL
Suppositories

If we were to use the superlatives which physicians all over the world have applied to Anusol Suppositories during the last twenty years, we would most certainly be accused of making exaggerated claims.

So we are not going to do it.

But we will gladly send you a sample.

And let you see what some physicians say about them.

MADE IN U.S.A.
FURNISHED TWELVE IN GREY BOX WHITE, RED AND BLACK LABEL AT BEFORE-THE-WAR PRICE.
SCHERING & GLATZ, INC. NEW YORK

Morse Wave Generator



The above apparatus is an instrument designed to synchronize normal muscle movements, producing definite wave muscle undulations, either for alternating or direct current effects. The former is exercise and massage and the latter to be used wherever the Galvanic current is indicated. The apparatus can be used on either the alternating or direct current without a transformer. The above features are characteristic of the apparatus.

MORSE-CHAPMAN COMPANY
462 Boylston Street, Boston, Mass.

VICTOR

Winning the Patient's Confidence

FEW patients have any real knowledge of medicine or surgery. Yet they form their own opinion of a practitioner's ability. How? By his manner, his skill, above all by the *character of his office and its equipment.*

The greater the physician's obvious technical resources, the greater must be the confidence that his patients have in him.

A modern office equipment convinces the patient that modern methods in diagnosing and treating disease are employed. It is not only a technical aid in practice. *It inspires confidence.*

No single piece of apparatus is of such general utility as a good X-ray equipment, and none that testifies more eloquently to the physician's resourcefulness. All practitioners recognize the diagnostic and therapeutic importance of the X-ray. Because he thinks that the taking of a radiograph requires special engineering knowledge, many a physician hesitates to install an X-ray equipment and thereby fails to satisfy his professional desire for thoroughness. The truth is that *the actual manipulation of the X-ray apparatus is as easily learned as that of the mechanical and electrical appliances with which most practitioners are familiar.*

The Victor X-Ray Corporation places its facilities and its long experience at the disposal of all practitioners. It will gladly send a technically informed representative to a physician who appreciates at its full worth the aid that the X-ray can lend him in his practice, but who wants engineering guidance. No obligation is incurred.

The physician who installs a Victor machine is entitled to receive Victor Service. This means that if his apparatus needs attention he does not have to communicate with the factory. The nearest Victor Service Station sends an expert, a man able to locate the source of trouble in a few moments. Victor Service is so organized that Victor apparatus need not be idle for many hours or days.

It is the purpose of Victor Service not only to help the physician in such emergencies, but also to give him mechanical and electrical guidance if he asks it, so that he may know how to secure the best results with his apparatus.

As part of this service policy the Victor X-Ray Corporation publishes a periodical called "Service Suggestions" in which X-ray progress is recorded primarily for the benefit of Victor clients. Others may find "Service Suggestions" of value. It will be sent to them on request.

Victor X-Ray Corporation

General Offices and Factory

Jackson Blvd. at Robey St. Chicago

Territorial Sales Distributors

Boston, Mass.
F. H. SAXBY & WESTON OYLER
711 Boylston Street

Oral Prophylaxis

MOUTH CLEANLINESS is no longer merely a matter of personal inclination or desire; medical men now recognize it as an essential detail in the prevention of disease.

THE MOUTH offers an ideal dwelling place for germs; it is of just the right temperature and moisture, and unless regularly cleansed, there are always enough decomposing food particles in the deep tissue folds and around the teeth to attract and favor the retention of all sorts of pathogenic bacteria.

NEGLECT OF THE MOUTH means a constant invitation to many of the acute diseases.

A CLEAN MOUTH is, therefore, the first consideration in the modern scheme of personal hygiene.

DIOXOGEN—the best and purest peroxide of hydrogen—is an invaluable mouth-cleanser. Owing its antiseptic qualities to pure oxygen—the same agent—that makes sunshine and good air the most efficient of germ-destroyers, Dioxogen penetrates to the deepest recesses of the mouth, where it softens and detaches all accumulated material, and cleanses and purifies everything it touches.

DIOXOGEN is not a mixture containing aromatic flavors or other substances to convey the idea of great antiseptic power. It is, instead, colorless, odorless and almost tasteless, with a germicidal efficiency superior to the solutions of carbolic or bichloride ordinarily used, but without the slightest toxic or irritating effect.

DIOXOGEN, in consequence, can be used as freely as necessary or desired, in even the youngest patients, with absolute certainty not only that it will keep the mouth and teeth in a clean and wholesome condition, but that it will always do its work in a pleasant and agreeable manner.

THE OAKLAND CHEMICAL COMPANY

59 Fourth Avenue
New York City



A useful booklet for the Roentgenologist

THIS little booklet should be in the hands of every Roentgenologist. It outlines in the following chapters, briefly, but practically, the methods which will promote more efficient use of X-Ray materials:

X-Ray Photographic Physics
The Generation of X-Rays, Exposure, Theory of Positioning, Radiation, Diaphragms, Screen Technique.
The Use of Dental X-Ray Films
Dark-Room Procedure
Developing Procedure
Developers and Fixing Baths
Lantern Slide Making and Copying.

Send for free copy.

EASTMAN KODAK CO., Rochester, N. Y.

Jump at Conclusions —False Diagnosis

The therapeutic value of pure petroleum oil has been established, in spite of the reaction resulting from over exploitation of and exaggerated claims for it.

Proper use of a perfected product assures satisfactory results.

TERRALINE

(Petroleum Purificatum)

a pioneer preparation of medicinally pure petroleum oil, bland, palatable and standardized, has for years demonstrated its value both as an intestinal lubricant and especially in the treatment of bronchial irritation, for the relief of cough, to promote expectoration and assist in the healing of inflamed areas.

Terraline is supplied either plain or with Creosote or Heroin. Terraline has been tested and proven by thousands of physicians, to whom it has appealed on account of its quality, its ethical introduction and its convincing response to the acid test of actual performance. Terraline forces conviction because it brings results.

Samples and literature to physicians only on request.

HILLSIDE CHEMICAL COMPANY

Newburgh, N. Y.

Pituitary Liquid

is the perfect preparation of Posterior Pituitary active principle. It, too, is without preservatives—½ c.c. obstetrical, 1 c.c. surgical.

Corpus Luteum

(*Armour*)

is true substance and will give results. Powder 2 and 5 gr. capsules and 2 and 5 gr. tablets.

Surgical Catgut Ligatures

Plain and chromic, regular (60 inch) emergency (20 inch) Iodized (60 inch)

Strong and sterile.

LABORATORY PRODUCTS

An Incomparable Product

The Suprarenalin preparations are now available.

Suprarenalin Powder . . . 1 grain vials
Suprarenalin Solution, 1:1000 . 1 ounce bottles
Suprarenalin Ointment, 1:1000 . . tubes

Suprarenalin designates the astringent, hemostatic and pressor principle of the Suprarenal Gland as isolated by the Armour chemists.

Suprarenalin Solution is the incomparable preparation of the kind. It is water-white, stable and non-irritating and is entirely free from chemical preservatives.

Suprarenalin ointment is bland and its effects very lasting.

ARMOUR AND COMPANY
CHICAGO

5821



A LARGE VARIETY OF MODELS, MAKING IT POSSIBLE TO SELECT THE PROPER PATTERN FOR ANY CONDITION REQUIRING ABDOMINAL SUPPORT. MADE TO ORDER AND CORRECTLY FITTED.



POMEROY COMPANY - 41 West St., Boston

SEPARATE DEPARTMENTS FOR MEN AND FOR WOMEN, WITH EXPERIENCED ATTENDANTS IN CHARGE. PHYSICIANS APPROVE OF OUR SERVICE BECAUSE IT INCLUDES FILLING PRESCRIPTIONS BY SUPPLYING THE KIND OF SUPPORT DESIRED.

WHEN NOT CONVENIENT TO SEND YOUR PATIENTS TO OUR FITTING ROOMS, WRITE FOR ORDER BLANKS AND DIRECTIONS FOR TAKING MEASUREMENTS.



PRESS OF JAMAICA PRINTING COMPANY, BOSTON, MASS.